


Dissecting a stellar explosion

April 3 2009



This artist's impression illustrates how a gamma-ray burst can flare dramatically over a short time period (gamma ray bursts usually last between a hundredth of a second to a hundred seconds). The bursts can occur as often as several times a day. There is no way to predict when or where they will next occur. ESA missions such as XMM-Newton, Integral and Ulysses study these mysterious, powerful bursts. Credits: ESA (Illustration by AOES Medialab)


(PhysOrg.com) -- Integral has captured one of the brightest gamma-ray bursts ever seen. A meticulous analysis of the data has allowed astronomers to investigate the initial phases of this giant stellar explosion, which led to the ejection of matter at velocities close to the speed of light. In particular, the astronomers believe that the explosion lifted a piece of the central engine's magnetic field into space.

On 19 December 2004, the blast from an exploding star arrived at Earth. ESA's Integral satellite, an orbiting gamma-ray observatory, recorded the entire event, providing information for what may prove to be one of the

most important gamma-ray bursts (GRBs) seen in recent years. As the data was collected, astronomers saw the 500-second-long burst rise to extraordinary brilliance.

"It is in the top 1% of the brightest GRBs we have seen," says Diego Götz, CEA Saclay, France, who headed the investigation.

The brightness of the event, known as GRB 041219A, has allowed the team to perform a painstaking investigation to extract a property known as the polarisation of the [gamma rays](#). The team have shown that the gamma rays were highly polarised and varied tremendously in level and orientation.



This artist's impression shows the centre of a dying star collapsing minutes before the star implodes. The blast from a Gamma Ray Burst is thought to be produced by a jet of fast-moving gas that bursts from near the central engine; probably a black hole created by such a collapse of the massive star. Credits: NASA/Dana Berr

Polarisation refers to the preferred direction in which the radiation wave oscillates. Polaroid sunglasses work with visible light by letting through only a single direction of polarisation, blocking most of the light from

entering our eyes.

The blast from a GRB is thought to be produced by a jet of fast-moving gas bursting from near the central engine; probably a black hole created by the collapse of the massive star. The polarisation is directly related to the structure of the [magnetic field](#) in the jet. So it is one of the best ways for astronomers to investigate how the central engine produces the jet. There are a number of ways this might happen.

In the first scenario, the jet carries a portion of the central engine's magnetic field into space. A second involves the jet generating the magnetic field far from the central engine. A third concerns the extreme case in which the jet contains no gas just magnetic energy, and a fourth scenario entails the jet moving through an existing field of radiation.

In each of the first three scenarios, the polarisation is generated by what is called synchrotron radiation. The magnetic field traps particles, known as electrons, and forces them to spiral, releasing polarised radiation. In the fourth scenario, the polarisation is imparted through interactions between the electrons in the jet and photons in the existing radiation field.

Götz believes that the Integral results favour a synchrotron model and, of those three, the most likely scenario is the first, in which the jet lifts the central engine's magnetic field into space. "It is the only simple way to do it," he says.

What Götz would most like to do is measure the polarisation for every GRB, to see whether the same mechanism applies to all. Unfortunately, many GRBs are too faint for the current instrumentation to succeed. Even the state-of-the-art IBIS instrument on Integral can only record the polarisation state of gamma rays if a celestial source is as bright as GRB 041219A.

"So, for now we just have to wait for the next big one," he says.

Source: European Space Agency ([news](#) : [web](#))

Citation: Dissecting a stellar explosion (2009, April 3) retrieved 3 May 2024 from
<https://phys.org/news/2009-04-stellar-explosion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.