

# Jeepers creepers: Climate change threatens endangered honeycreepers

May 26 2009

---

As climate change causes temperatures to increase in Hawaii's mountains, deadly non-native bird diseases will likely also creep up the mountains, invading most of the last disease-free refuges for honeycreepers - a group of endangered and remarkable birds.

A just-published U.S. Geological Survey (USGS) review discusses the likelihood of a forthcoming "disease invasion" by examining the present altitudinal range of avian malaria and pox, honeycreeper distribution, and the future projected range of diseases and honeycreeper habitat with [climate change](#).

At one time, the Hawaiian Islands had no mosquitoes - and no mosquito-borne diseases. But, by the late 1800s, mosquitoes had set up permanent housekeeping, setting the stage for epidemic transmission of avian malaria and pox. Honeycreepers - just like people faced with novel viruses such as [swine flu](#) - had no natural resistance against these diseases.

Before long, Hawaii's native honeycreepers significantly declined in numbers and geographic range. It was likely that malaria swept rapidly across all of the lower [Hawaiian Islands](#) after the disease was introduced, leaving few survivors. Today, native Hawaiian birds face one of the highest rates of extinction in the world. Of 41 honeycreeper species and subspecies known since historic times, 17 are probably extinct, 14 are endangered, and only 3 are in decent shape.

Pox and malaria transmission in Hawaii depends on climatic conditions, especially seasonal changes in temperature and rainfall that increase or decrease mosquito populations. "Without question, the one factor that prevented widespread and rapid extinction of virtually all of Hawaii's native honeycreepers after the introduction of avian pox and avian malaria was the presence of high-altitude disease refuges on Kauai, Maui and Hawaii," said lead study author Dr. Carter Atkinson, a USGS microbiologist based at the USGS Pacific Islands Ecosystems Research Center in Hawaii.

These cool, high-elevation - above 4,000 feet (1,200 meters) -- mountains not only provided habitats that mosquitoes didn't thrive in, but they also had habitat that honeycreepers liked, wrote the authors. While birds in those areas find refuge from the diseases - dispersing juvenile birds and adults that follow seasonal flowering of native plants to lower elevations are exposed to disease.

"Unfortunately," said study co-author, USGS scientist Dr. Dennis LaPointe, "this seasonal movement happens at the same time that mosquito populations soar at mid-elevations, which fuels high disease-transmission rates there. There's a continuous source of disease-susceptible birds each fall."

Although most disease transmission now occurs in these mid-elevation forests, this will change if the projected 3.6 degrees Fahrenheit (2 degrees Centigrade) raise in temperature occurs.

"With this kind of temperature change, about 60 to 96 percent of the high-elevation disease refuges would disappear," said Atkinson. For example, available high-elevation forest habitat in the low-risk disease zone would likely decline by nearly 60 percent at Hanawi Natural Area Reserve on Maui to as much as 96 percent at Hakalau Forest National Wildlife Refuge on Hawaii Island. On other islands, such as Kauai, with

lower elevations and no low-risk zones even now, predicted temperature changes would likely be catastrophic for remaining honeycreeper species.

"Right now, disease transmission in the mountains of Kauai is highly seasonal, but with temperature increases, disease would be able to be transmitted throughout most of the year," said Atkinson.

In addition, the tropical inversion layer - often visible as a thin cloud layer around high peaks -- may play a more significant role than temperature in determining tree line and the upper extent of forest bird habitat, the authors wrote. The inversion layer forms as cool, dry air meets warm, moist air, creating an inversion layer that caps moisture and cloud development between 5,900 and 7,900 feet (1,800 and 2,400 meters). Over the past 24 years, the height of this layer has remained fairly stable in spite of warming trends but has increased in how often it occurs.

"This could have a tremendous adverse effect on the continued existence of high-elevation disease refugia," said Atkinson. "Remaining bird populations could be squeezed between expanding disease transmission from lower elevations and the upper limits of suitable forest habitat. Such changes would likely push remaining populations of threatened and endangered honeycreepers to extinction, and cause severe declines in other honeycreepers not now endangered but susceptible to avian malaria."

Given the likelihood of global warming, the authors suggested that management of mid-elevation habitats to reduce disease transmission will become increasingly vital. The best opportunities for doing this, they wrote, will be through reducing habitat for mosquito larvae, habitat that is often created by introduced feral pigs, goats and cattle in forests.

"The survival of these species into the next century may ultimately depend on our ability to remove or offset introduced threats and restore native forests from sea level to tree line," Atkinson said.

Honeycreepers rival Darwin's finches in the Galapagos Islands in terms of their bill types and number of species that descended from a common founder. The birds specialize on food that includes nectar, fruits and insects. Before people came to the islands, as many as 56 kinds of honeycreepers probably occurred.

Avian malaria is caused by a protozoan parasite, and avian pox is a viral infection that typically causes tumor-like swellings on exposed skin of the feet, legs, beak and eyelids of infected birds. Malaria often results in appetite and weight loss, anemia, and massive enlargement of birds' liver and spleen.

More information: The article, Introduced avian diseases, climate change, and the future of Hawaiian honeycreepers, was published in the *Journal of Avian Medicine and Surgery*.

Source: United States Geological Survey ([news](#) : [web](#))

Citation: Jeepers creepers: Climate change threatens endangered honeycreepers (2009, May 26) retrieved 1 May 2024 from

<https://phys.org/news/2009-05-jeepers-creepers-climate-threatens-endangered.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.