

## **Evolution can occur in less than 10 years**

June 11 2009



Guppies are small fresh-water fish that biologists have studied for long. Credit: Paul Bentzen.

How fast can evolution take place? In just a few years, according to a new study on guppies led by UC Riverside's Swanne Gordon, a graduate student in biology.

Gordon and her colleagues studied <u>guppies</u> — small fresh-water fish biologists have studied for long — from the Yarra River, Trinidad. They introduced the guppies into the nearby Damier River, in a section above a barrier waterfall that excluded all predators. The guppies and their descendents also colonized the lower portion of the stream, below the barrier waterfall, that contained natural predators.

Eight years later (less than 30 guppy generations), the researchers found



that the guppies in the low-predation environment above the barrier waterfall had adapted to their new environment by producing larger and fewer offspring with each reproductive cycle. No such adaptation was seen in the guppies that colonized the high-predation environment below the barrier waterfall.

"High-predation females invest more resources into current reproduction because a high rate of mortality, driven by predators, means these females may not get another chance to reproduce," explained Gordon, who works in the lab of David Reznick, a professor of biology. "Lowpredation females, on the other hand, produce larger embryos because the larger babies are more competitive in the resource-limited environments typical of low-predation sites. Moreover, low-predation females produce fewer embryos not only because they have larger embryos but also because they invest fewer resources in current reproduction."

Study results appear in the July issue of The American Naturalist.

Natural guppy populations can be divided into two basic types. Highpredation populations are usually found in the downstream reaches of rivers, where they coexist with predatory fishes that have strong effects on guppy demographics. Low-predation populations are typically found in upstream tributaries above barrier waterfalls, where strong predatory fishes are absent. Researchers have found that this broad contrast in predation regime has driven the evolution of many adaptive differences between the two guppy types in color, morphology, behavior, and life history.

Gordon's research team performed a second experiment to measure how well adapted to survival the new population of guppies were. To this end, they introduced two new sets of guppies, one from a portion of the Yarra River that contained predators and one from a predator-free tributary to



the Yarra River into the high-and low-predation environments in the Damier River.

They found that the resident, locally adapted guppies were significantly more likely to survive a four-week time period than the guppies from the two sites on the Yarra River. This was especially true for juveniles. The adapted population of juveniles showed a 54-59 percent increase in survival rate compared to their counterparts from the newly introduced group.

"This shows that adaptive change can improve survival rates after fewer than ten years in a new environment," Gordon said. "It shows, too, that evolution might sometimes influence population dynamics in the face of environmental change."

Source: University of California - Riverside (<u>news</u> : <u>web</u>)

Citation: Evolution can occur in less than 10 years (2009, June 11) retrieved 10 May 2024 from <u>https://phys.org/news/2009-06-evolution-years.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.