


Fermilab's CDF observes Omega-sub-b baryon

June 29 2009



Six quarks--up, down, strange, charm, bottom and top--are the building blocks of matter. Protons and neutrons are made of up and down quarks, held together by the strong nuclear force. The CDF experiment now has observed the Omega-sub-b particle, which contains two strange quarks (s) and one bottom quark (b).


(PhysOrg.com) -- At a recent physics seminar at the Department of Energy's Fermi National Accelerator Laboratory, Fermilab physicist Pat Lukens of the CDF experiment announced the observation of a new particle, the Omega-sub-b (Ω_b). The particle contains three quarks, two

strange quarks and a bottom quark (s-s-b). It is an exotic relative of the much more common proton and has about six times the proton's mass.

The observation of this “doubly strange” particle, predicted by the Standard Model, is significant because it strengthens physicists’ confidence in their understanding of how quarks form matter. In addition, it conflicts with a 2008 result announced by CDF’s sister experiment, DZero.

The Omega-sub-b is the latest entry in the "periodic table of baryons." Baryons are [particles](#) formed of three quarks, the most common examples being the proton and neutron. The Tevatron particle accelerator at Fermilab is unique in its ability to produce baryons containing the b quark, and the large data samples now available after many years of successful running enable experimenters to find and study these rare particles. The observation opens a new window for scientists to investigate its properties and better understand this rare object.

Combing through almost half a quadrillion (1000 billion) proton-antiproton collisions produced by Fermilab's Tevatron particle collider, the CDF collaboration isolated 16 examples in which the particles emerging from a collision revealed the distinctive signature of the Omega-sub-b. Once produced, the Omega-sub-b travels a fraction of a millimeter before it decays into lighter particles. This decay, mediated by the weak force, occurs in about a trillionth of a second. In fact, CDF has performed the first ever measurement of the Omega-sub-b lifetime and obtained $1.13 +0.53-0.40$ (stat.) ± 0.02 (syst.) trillionths of a second.



Baryons are particles made of three quarks. The quark model predicts the combinations that exist with either spin $J=1/2$ (this graphic) or spin $J=3/2$. The graphic shows the various three-quark combinations with $J=1/2$ that are possible using the three lightest quarks--up, down and strange--and the bottom quark. The CDF collaboration observed the Omega-sub-b, highlighted in the graphic. There exist additional baryons involving the charm quark, which are not shown. The top quark, discovered at Fermilab in 1995, is too short-lived to become part of a baryon.

In August 2008, the DZero experiment announced its own observation of the Omega-sub-b based on a smaller sample of Tevatron data.

Interestingly, the new CDF observation announced here is in direct conflict with the earlier DZero result. The CDF physicists measured the Omega-sub-b mass to be $6054.4 \pm 6.8(\text{stat.}) \pm 0.9(\text{syst.}) \text{ MeV}/c^2$, compared to DZero's $6165 \pm 10(\text{stat.}) \pm 13(\text{syst.}) \text{ MeV}/c^2$. These two experimental results are statistically inconsistent with each other leaving scientists from both experiments wondering whether they are measuring the same particle. Furthermore, the experiments observed different rates of production of this particle. Perhaps most interesting is that neither experiment sees a hint of evidence for the particle at the other's measured value.

Although the latest result announced by CDF agrees with theoretical expectation for the Omega-sub-b both in the measured production rate and in the mass value, further investigation is needed to solve the puzzle of these conflicting results.

The Omega-sub-b discovery follows the observation of the Cascade-b-minus baryon (Ξ_b), first observed at the Tevatron in 2007, and two types of Sigma-sub-b baryons (Σ_b), discovered at the Tevatron in 2006.

The CDF collaboration submitted a paper that summarizes the details of its discovery to the journal Physical Review D. It is available online at: arxiv.org/abs/0905.3123

Provided by [Fermilab](#) ([news](#) : [web](#))

Citation: Fermilab's CDF observes Omega-sub-b baryon (2009, June 29) retrieved 3 May 2024 from <https://phys.org/news/2009-06-fermilab-cdf-omega-sub-b-baryon.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--