


NIST researchers 'all aglow' over new test of toxin strength

June 17 2009



Microscopic photographs showing the level of fluorescence (green images on left) and the numbers of cells (blue images on right) at the time the cells were exposed to ricin (top row) and six hours later (bottom row). Note that the fluorescence has decreased (more black regions in the bottom left photo) while the cell count remains virtually the same. This shows that the NIST assay detects ricin even before significant cell death occurs. Credit: M. Halter, NIST

A new National Institute of Standards and Technology (NIST) assay using a “glow or no glow” technique may soon help the U.S. Department of Homeland Security (DHS) defend the nation against a spectrum of biological weapons that could be used in a terrorist attack. One very dangerous toxin on the list is ricin, a protein derived from castor beans that is lethal in doses as small as 500 micrograms—about the size of a grain of salt.

As part of its efforts to address the threat, DHS is working with NIST to create a standardized ricin sample with a known potency. Such a standard is needed both to check the accuracy of detection equipment, and, should an attack occur, to confirm the success of decontamination procedures. A major step toward this goal—the development of a rapid, reliable and precise assay of the potency of a ricin sample—has now been achieved by NIST scientists.

As detailed in an article posted online this week in *Assay and Drug Development Technologies*, the new NIST assay uses a commercially available cell line genetically engineered to produce large amounts of green fluorescent protein (GFP). Ricin shuts down ribosomes—the protein manufacturing factories of a cell. Assay cells exposed to the toxin will quickly stop synthesizing GFP. This, in turn, will result in a measurable decrease in fluorescence—a drop that correlates directly to the strength of the ricin present.

The NIST assay yields many advantages over traditional cytotoxicity measuring systems, including: a highly sensitive response to ricin (as little as 1 nanogram per milliliter) within six rather than 24 hours; detection of the toxin long before significant cell death has occurred; results that are highly reproducible; no need for the addition of any reagents; and the flexibility to measure the potency of other ribosome inhibitors, even nanoparticles, with the same precision as ricin.

Citation: M. Halter, J.L. Almeida, A. Tona, K.D. Cole, A.L. Plant and J.T. Elliott. A mechanistically relevant cytotoxicity assay based on the detection of cellular green fluorescent protein. *Assay and Drug Development Technologies*, Vol. 7, No. 4 (August 2009; posted online the week of June 15, 2009).

Source: National Institute of Standards and Technology (NIST)

Citation: NIST researchers 'all aglow' over new test of toxin strength (2009, June 17) retrieved 27 April 2024 from <https://phys.org/news/2009-06-nist-aglow-toxin-strength.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.