


Physicists Propose a Method to Observe Dirac Monopoles

July 28 2009, By Lisa Zyga



Possible configurations for the spin texture of a Bose-Einstein condensate. These defects give rise to a vorticity field that is equivalent to the magnetic field of a magnetic monopole. Credit: Pietilä and Möttönen.

(PhysOrg.com) -- For decades, scientists have been intrigued by the hypothetical existence of magnetic monopoles - particles with only a north or south magnetic pole, thus having a nonzero magnetic charge. But while modern theories predict their existence, monopoles have not yet been experimentally observed, despite numerous attempts. Recently, physicists have introduced a new method to create analogies to magnetic monopoles, and predict that they could be observed with current technology.

Ville Pietilä and Mikko Möttönen, both of the Helsinki University of Technology in Finland and the University of New South Wales in

Australia, have published their theoretical demonstration in a recent issue of *Physical Review Letters*. Here, they explain how applying an external [magnetic field](#) to a Bose-Einstein condensate (BEC) - a large group of cold atoms that exhibits coherent quantum properties - can create point-like topological defects on the spin texture of the BEC. These defects give rise to a vorticity field that is essentially equivalent to the magnetic field of a magnetic monopole.

“Since all experimental attempts to find magnetic monopoles have proven to be futile, there is no experimental evidence supporting the existence of magnetic monopoles,” Pietilä told *PhysOrg.com*. “Other types of monopoles without the Dirac string have been realized in experiments already in the early ‘90s in liquid crystals. An analogy of the real space magnetic monopole was reported to occur in the crystal momentum space of a metallic ferromagnet, but the experimental evidence in this case is somewhat indirect. Dirac monopoles in the more general settings are predicted to occur in various systems such as superfluid Helium-3 and dilute spin-1 Bose-Einstein condensate but so far there are no (direct) experimental observations although they may have been present in some of the Helium-3 experiments. There is also a very recent suggestion on how to induce a magnetic monopole to a band insulator.

“Since magnetic monopoles have never been observed, it is pertinent to ask whether there is something unphysical in the whole concept,” he continued. “Our work shows that at least the Dirac monopole can be realized experimentally, thus indicating that it is more than just a theoretical curiosity. However, it should be stressed that our work does not tell anything about the existence of magnetic monopoles in the electromagnetic field.”

Pietilä and Möttönen predict that it should be possible to design an experiment to detect the monopole in this situation, if it does exist. As

they explain, the magnetic field of the monopole is provided by a Dirac string, which is a line extending from the monopole to infinity. The Dirac string explains why the monopole charge comes in discrete quanta. Since the Dirac string carries two quanta of angular momentum, it is expected to be prone to splitting into two separate strings, each carrying a single quantum.

“The Dirac monopole is conceptually the simplest model for a magnetic point charge (i.e. a magnetic monopole) and this makes the Dirac monopole important,” Pietilä said. “Monopoles in general can be any point-like defects in a three-component vector field defined in three spatial dimensions, but they are not necessarily associated with a singular filament (the Dirac string).”

Through modeling, Pietilä and Möttönen theoretically demonstrate that the Dirac string can be split into two parts by modifying the external magnetic field, which breaks the rotation symmetry of the system. In this case, state-of-the-art experimental methods that measure changes in particle density and spin density could be used to detect a signature of the monopole.

In addition, in spin-1 BEC, the spin texture is topologically unstable, allowing it to be removed. Through simulations, the scientists show that removing the monopole could be done by turning off the external magnetic fields. The monopole “unwinds” itself along the Dirac string and leaves behind a closed vortex ring. The speed of the unwinding depends on how quickly or gradually the magnetic field is turned off.

The existence of magnetic monopoles has far-reaching implications for many research areas, including the laws of quantum mechanics, theories of elementary [particles](#), and cosmology. Experimentally confirming the existence of monopoles would provide long-sought evidence for some ideas, and possibly open up the doors to other new ones.

“The existence of magnetic monopoles implies that the electric charge has to be quantized (that is, it can only appear as an integral multiple of the elementary charge),” Pietilä said. “Existence of magnetic monopoles has also certain cosmological implications. Since monopoles in general are related to the spontaneous symmetry breaking, they impact the unified theories describing the interactions between the elementary particles. In the context of condensed matter physics, monopoles typically arise as a manifestation of some exotic collective behavior of matter and can be used to explain the unusual characteristics of such systems.”

More information: Ville Pietilä and Mikko Möttönen. “Creation of Dirac Monopoles in Spinor Bose-Einstein Condensates.” *Physical Review Letters* 103, 030401 (2009).

link.aps.org/doi/10.1103/PhysRevLett.103.030401

Copyright 2009 PhysOrg.com.

All rights reserved. This material may not be published, broadcast, rewritten or redistributed in whole or part without the express written permission of PhysOrg.com.

Citation: Physicists Propose a Method to Observe Dirac Monopoles (2009, July 28) retrieved 20 April 2024 from <https://phys.org/news/2009-07-physicists-method-dirac-monopoles.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--