

Helping Albacore tuna come out of the can

August 11 2009



New information on the biology of albacore tuna will assist in stock assessments and providing advice on harvest levels in Australian and international waters.

Photo by: CSIRO/FRDC

(PhysOrg.com) -- Scientists are set to study the earbones and organs of more than 2000 albacore tuna to better understand the growth, age and breeding patterns of this increasingly important species.

Albacore tuna are harvested from tropical to temperate waters, mainly for canning, but also to satisfy the growing market for fresh [fish](#) in Europe and the United States.

While previously caught as by-product, albacore have become a prime target of Australia's Eastern Tuna and Billfish Fishery which operates from Cape York to Tasmania and on the adjacent high seas. Several Pacific island nations and distant water fleets, including Taiwan and Korea, also target albacore.

This rising catch of albacore has generated a need for more information about the species to assess stock levels and provide advice on harvest levels in Australian and international waters.

To meet this need, scientists from the CSIRO Wealth from Oceans Flagship are studying the biology of albacore stocks from eastern Australia and the wider south-western Pacific Ocean to determine their age, growth and reproductive characteristics.

The two-year study - funded by the Fisheries Research and Development Corporation and CSIRO, the Department of Agriculture, Fisheries and Forestry, and the Western and Central Pacific [Fisheries](#) Commission - builds on earlier studies in 2007 and 2008-09.

Project leader, Jess Farley, says fishery-wide information about how albacore age, grow and reproduce is needed to provide reliable stock assessments and set appropriate levels for future catches.

The age of the albacore will be estimated by interpreting growth bands formed in their earbones (otoliths) and dorsal spines in a process similar to reading growth rings of trees. Albacore reproductive organs will be examined to determine the location, timing, frequency and magnitude of spawning.

“Previous studies have shown that albacore grow to 1.2 metres in length and at least 30 kilograms in weight, and live up to 14 years,” Ms Farley says.

“More detailed biological information will allow us to calculate growth rates, age at maturity and longevity, and to see whether these differ by gender and region.

“We aim to develop a ‘maturity schedule’ for female albacore across the

south Pacific to help estimate their spawning output at different sizes and ages. This is the key to understanding the productivity of the stocks.”

CSIRO scientist Dr Campbell Davies says that, like other fish species, tunas release many thousands, if not millions, of eggs in a spawning season and individual spawning output increases with the size and age of the female fish.

“This needs to be quantified so that stock-assessment scientists can estimate how different levels of fishing will affect the spawning capacity of the whole population,” he says.

The CSIRO scientists will collaborate with the Secretariat of the Pacific Community (SPC) on the ageing and reproductive work, which will link with an SPC tagging program examining albacore movements in the Western and Central Pacific Ocean. The fishing industry will provide albacore samples and an archival reference collection will be established.

Provided by CSIRO ([news](#) : [web](#))

Citation: Helping Albacore tuna come out of the can (2009, August 11) retrieved 19 April 2024 from <https://phys.org/news/2009-08-albacore-tuna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.