

# Researchers show how organic carbon compounds emitted by trees affect air quality

August 6 2009

---



Caltech scientists used mass spectrometers on NASA's DC8 aircraft to measure epoxides above Canada's Boreal Forest last summer. These epoxides are formed from isoprene, a chemical emitted into the atmosphere by many deciduous plants. Credit: Yohei Shinozuka

A previously unrecognized player in the process by which gases produced by trees and other plants become aerosols—microscopically small particles in the atmosphere—has been discovered by a research team led by scientists at the California Institute of Technology.

Their research on the creation and effects of these chemicals, called epoxides, is being featured in this week's issue of the journal *Science*.

Paul Wennberg, the R. Stanton Avery Professor of [Atmospheric](#)

[Chemistry](#) and Environmental Science and Engineering and director of the Ronald and Maxine Linde Center for Global Environmental Science at Caltech, and John Seinfeld, the Louis E. Nohl Professor and professor of chemical engineering, have been studying the role of biogenic emissions—organic carbon compounds given off by plants and trees—in the atmospheric chemical reactions that result in the creation of aerosols.

"If you mix emissions from the city with emissions from plants, they interact to alter the chemistry of the atmosphere," Wennberg notes.

While there's been plenty of attention paid to the effect of emissions from cars and manufacturing, less is understood about what happens to biogenic emissions, especially in places where there are relatively few man-made emissions. That situation is the focus of the research that led to this Science paper. "What we're interested in," Wennberg explains, "is what happens to the chemicals produced by trees once they are emitted into the atmosphere."

In these studies, the research team focused on a chemical called isoprene, which is given off by many deciduous trees. "The king emitters are oaks," Wennberg says. "And the isoprene they emit is one of the reasons that the Smoky Mountains appear smoky."

Isoprene is no minor player in atmospheric chemistry, Wennberg notes. "There is much more isoprene emitted to the atmosphere than all of the gases—gasoline, industrial chemicals—emitted by human activities, with the important exceptions of [methane](#) and carbon dioxide," he says. "And isoprene only comes from plants. They make hundreds of millions of tons of this chemical . . . for reasons that we still do not fully understand."

"Much of the emission of isoprene occurs where anthropogenic emissions are limited," adds Caltech graduate student Fabien Paulot, the

paper's first author. "The chemistry is very poorly understood."

Once released into the atmosphere, isoprene gets "oxidized or chewed on" by free-radical oxidants such as OH, explains Wennberg. It is this chemistry that is the focus of this new study. In particular, the research was initiated to understand how the oxidation of isoprene can lead to formation of atmospheric particulate matter, so-called secondary organic aerosol. "A small fraction of the isoprene becomes secondary organic aerosol," Seinfeld notes, "but because isoprene emissions are so large, even this small fraction is important."

Up until now, the chemical pathways from isoprene to aerosol were not known. Wennberg, Seinfeld, and their colleagues discovered that this aerosol likely forms from chemicals known as an epoxides.

The name is apt. "These epoxides are nature's glue," says Wennberg. And, much like the epoxy you buy in a hardware store—which requires the addition of an acid for the compound to turn into glue—the epoxides found in the atmosphere also need an acidic kick in order to become sticky.

"When these epoxides bump into particles that are acidic, they make glue," Wennberg explains. "The epoxides precipitate out of the atmosphere and stick to the particles, growing them and resulting in lowered visibility in the atmosphere." Because the acidity of the aerosols is generally higher in the presence of anthropogenic activities, the efficiency of converting the epoxides to aerosol is likely higher in polluted environments, illustrating yet another complex interaction between emissions from the biosphere and from humans.

"Particles in the atmosphere have been shown to impact human health, as they are small enough to penetrate deep into the lungs of people. Also, [aerosols](#) impact Earth's climate through the scattering and absorption of

solar radiation and through serving as the nuclei on which clouds form. So it is important to know where particles come from," notes Seinfeld.

The research team was able to make this scientific leap forward thanks to their development of a new type of chemical ionization mass spectrometry (CIMS), led by coauthor and Caltech graduate student John Crouse. "These new CIMS methods open up a very wide range of possibilities for the study of new sets of compounds that scientists have been largely unable to measure previously, mainly because they decompose when analyzed with traditional techniques."

In general, molecules identified and quantified using mass spectroscopy must first be converted to charged ions. They are then directed into an electric field, where the ions are sorted by mass. The problem with traditional ionization techniques is that delicate molecules, such as those produced in the oxidation of isoprene, generally fragment during the ionization process, making their identification difficult or impossible. "This new method was originally developed in order to allow scientists to make atmospheric measurements from airplanes. It is able to ionize gasses, even fragile peroxide compounds, while still preserving information about the size or mass of the original molecule," says Wennberg.

That makes determining the individual gases in a complex mixture much easier—especially when, as it turned out, you're looking at a chemical you weren't expecting to find.

Wennberg and colleagues also used oxygen isotopes—oxygen atoms with different numbers of neutrons in their nucleus, and thus different masses—to gain insight into the chemical mechanism yielding epoxides. Epoxides have remained unidentified so far because they have the same mass as another chemical that had been anticipated to form in isoprene oxidation, peroxide. "The oxygen isotopes separated the peroxides from

epoxides and further showed that as the epoxides form, OH is recycled to the atmosphere," comments Paulot. "Since OH is the atmosphere detergent, cleaning the atmosphere of many chemicals, the recycling has important implications for the overall oxidizing capacity of the atmosphere."

The identification of a major photochemical pathway to formation of epoxides helps to explain just how tree emissions of organic carbon compounds influence the air in both city and rural settings. While trees aren't exactly the "killers" that Ronald Reagan was once so famously derided for calling them, their isoprene emission levels can—and often probably should—"be a part of the criteria we use when buying and planting trees in a polluted urban setting," notes Wennberg. In fact, he points out, the South Coast Air Quality Management District in Southern California already does this with its list of "approved" trees that don't emit large amounts of organic carbon compounds into the atmosphere.

Source: California Institute of Technology ([news](#) : [web](#))

Citation: Researchers show how organic carbon compounds emitted by trees affect air quality (2009, August 6) retrieved 11 September 2024 from <https://phys.org/news/2009-08-carbon-compounds-emitted-trees-affect.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--