


New nanolaser -- spaser -- key to future optical computers and technologies

August 16 2009, by Emil Venere



Researchers have created the tiniest laser since its invention nearly 50 years ago. Because the new device, called a "spaser," is the first of its kind to emit visible light, it represents a critical component for possible future technologies based on "nanophotonic" circuitry. The color diagram (a) shows the nanolaser's design: a gold core surrounded by a glasslike shell filled with green dye. Scanning electron microscope images (b and c) show that the gold core and the thickness of the silica shell were about 14 nanometers and 15 nanometers, respectively. A simulation of the SPASER (d) shows the device emitting visible light with a wavelength of 525 nanometers. (Birck Nanotechnology Center, Purdue University)

(PhysOrg.com) -- Because the new device, called a "spaser," is the first of its kind to emit visible light, it represents a critical component for possible future technologies based on "nanophotonic" circuitry, said Vladimir Shalaev, the Robert and Anne Burnett Professor of Electrical and Computer Engineering at Purdue University.

Such circuits will require a laser-light source, but current lasers can't be made small enough to integrate them into electronic chips. Now researchers have overcome this obstacle, harnessing clouds of electrons

called "surface plasmons," instead of the photons that make up light, to create the tiny spasers.

Findings are detailed in a paper appearing online Sunday (Aug. 16) in the journal *Nature*, reporting on work conducted by researchers at Purdue, Norfolk State University and Cornell University.

Nanophotonics may usher in a host of radical advances, including powerful "hyperlenses" resulting in sensors and microscopes 10 times more powerful than today's and able to see objects as small as DNA; computers and consumer electronics that use light instead of electronic signals to process information; and more efficient solar collectors.

"Here, we have demonstrated the feasibility of the most critical component - the nanolaser - essential for nanophotonics to become a practical technology," Shalaev said.

The "spaser-based nanolasers" created in the research were spheres 44 [nanometers](#), or billionths of a meter, in diameter - more than 1 million could fit inside a [red blood cell](#). The spheres were fabricated at Cornell, with Norfolk State and Purdue performing the optical characterization needed to determine whether the devices behave as lasers.

The findings confirm work by physicists David Bergman at Tel Aviv University and Mark Stockman at Georgia State University, who first proposed the spaser concept in 2003.

"This work represents an important milestone that may prove to be the start of a revolution in [nanophotonics](#), with applications in imaging and sensing at a scale that is much smaller than the wavelength of visible light," said Timothy D. Sands, the Mary Jo and Robert L. Kirk Director of the Birck Nanotechnology Center in Purdue's Discovery Park.

The spasers contain a gold core surrounded by a glasslike shell filled with green dye. When a light was shined on the spheres, plasmons generated by the gold core were amplified by the dye. The plasmons were then converted to photons of [visible light](#), which was emitted as a [laser](#).

Spaser stands for surface plasmon amplification by stimulated emission of radiation. To act like lasers, they require a "feedback system" that causes the surface plasmons to oscillate back and forth so that they gain power and can be emitted as light. Conventional lasers are limited in how small they can be made because this feedback component for photons, called an optical resonator, must be at least half the size of the wavelength of [laser light](#).

The researchers, however, have overcome this hurdle by using not photons but surface plasmons, which enabled them to create a resonator 44 nanometers in diameter, or less than one-tenth the size of the 530-nanometer wavelength emitted by the spaser.

"It's fitting that we have realized a breakthrough in laser technology as we are getting ready to celebrate the 50th anniversary of the invention of the laser," Shalaev said.

The first working laser was demonstrated in 1960.

The research was conducted by Norfolk State researchers Mikhail A. Noginov, Guohua Zhu and Akeisha M. Belgrave; Purdue researchers Reuben M. Bakker, Shalaev and Evgenii E. Narimanov; and Cornell researchers Samantha Stout, Erik Herz, Teeraporn Suteewong and Ulrich B. Wiesner.

Future work may involve creating a spaser-based nanolaser that uses an electrical source instead of a light source, which would make them more

practical for computer and electronics applications.

Source: Purdue University ([news](#) : [web](#))

Citation: New nanolaser -- spaser -- key to future optical computers and technologies (2009, August 16) retrieved 25 April 2024 from <https://phys.org/news/2009-08-nanolaser-spaser-key-future.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.