

# Trigger-Happy Star Formation

August 12 2009

---



This composite image, combining data from the Chandra X-ray Observatory and the Spitzer Space Telescope shows the molecular cloud Cepheus B, located in our Galaxy about 2,400 light years from the Earth. A molecular cloud is a region containing cool interstellar gas and dust left over from the formation of the galaxy and mostly contains molecular hydrogen. The Spitzer data, in red, green and blue shows the molecular cloud (in the bottom part of the image) plus young stars in and around Cepheus B, and the Chandra data in violet shows the young stars in the field. Credits: X-ray: NASA/CXC/PSU/K. Getman et al.; IRL NASA/JPL-Caltech/CfA/J. Wang et al.

(PhysOrg.com) -- A new study from two of NASA's Great Observatories provides fresh insight into how some stars are born, along with a

beautiful new image of a stellar nursery in our Galaxy. The research shows that radiation from massive stars may trigger the formation of many more stars than previously thought.

While astronomers have long understood that [stars](#) and planets form from the collapse of a cloud of gas, the question of the main causes of this process has remained open.

One option is that the cloud cools, gravity gets the upper hand, and the cloud falls in on itself. The other possibility is that a "trigger" from some external source -- like radiation from a massive star or a shock from a [supernova](#) -- initiates the collapse. Some previous studies have noted a combination of triggering mechanisms in effect.

By combining observations of Cepheus B from the [Chandra X-ray Observatory](#) and [Spitzer Space Telescope](#), researchers have taken an important step in addressing this question. Cepheus B is a cloud of mainly cool molecular hydrogen located about 2,400 light years from the Earth. There are hundreds of very young stars inside and around the cloud -- ranging from a few millions years old outside the cloud to less than a million in the interior -- making it an important testing ground for [star formation](#).

"Astronomers have generally believed that it's somewhat rare for stars and planets to be triggered into formation by radiation from massive stars," said Konstantin Getman of Penn State University, and lead author of the study. "Our new result shows this belief is likely to be wrong."

This particular type of triggered star formation had previously been seen in small populations of a few dozen stars, but the latest result is the first time it has been clearly observed in a rich population of several hundred stars.

While slightly farther away than the famous Orion star-forming region, Cepheus B is at a better orientation for astronomers to observe the triggering process. The Chandra observations allowed the astronomers to pick out young stars within and around Cepheus B. Young stars have turbulent interiors that generate highly active magnetic fields, which, in turn, produce strong and identifiable X-ray signatures.

The Spitzer data revealed whether the young stars have a disk of material (known as "protoplanetary" disks) around them. Since they only exist in very young systems where planets are still forming, the presence of protoplanetary disks -- or lack thereof -- is an indication of the age of a star system.

The new study suggests that star formation in Cepheus B is mainly triggered by radiation from one bright, massive star outside the molecular cloud. According to theoretical models, radiation from this star would drive a compression wave into the cloud triggering star formation in the interior, while evaporating the cloud's outer layers. The Chandra-Spitzer analysis revealed slightly older stars outside the cloud while the youngest stars with the most protoplanetary disks congregate in the cloud interior -- exactly what is predicted from the triggered star formation scenario.

"We essentially see a wave of star and planet formation that is rippling through this cloud," said co-author Eric Feigelson, also of Penn State. "Outside the cloud, the stars probably have newly born planets while inside the cloud the planets are still gestating."

Previous observations of Cepheus B had shown a rim of ionized gas around the molecular cloud and facing the massive star. However, the wave of star formation -- an additional crucial feature to identifying the source of the star formation -- had not previously been seen. "We can even clock how quickly this wave is traveling and it's going about 2,000

miles per hour," said Getman.

The star that is the catalyst for the star formation in Cepheus B, is about 20 times as massive as the Sun, or at least five times weightier than any of the other stars in Cepheus B.

The Chandra and Spitzer data also suggest that multiple episodes of star and planet formation have occurred in Cepheus B over millions of years and that most of the material in the cloud has likely already been evaporated or transformed into stars.

"It seems like this nearby cloud has already made most of its stars and its fertility will soon wane," said Feigelson. "It's clear that we can learn a lot about stellar nurseries by combining data from these two Great Observatories."

A paper describing these results was published in the July 10 issue of the *Astrophysical Journal*. The team of astronomers that worked with Getman and Feigelson also included Kevin Luhman and Gordon Garmire from Penn State, Aurora Sicilia-Aguilar from Max-Planck-Institut für Astronomie, and Junfeng Wang from Harvard-Smithsonian Center for Astrophysics.

Provided by JPL/NASA ([news](#) : [web](#))

Citation: Trigger-Happy Star Formation (2009, August 12) retrieved 1 May 2024 from <https://phys.org/news/2009-08-trigger-happy-star-formation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--