

# Research team finds first evolutionary branching for bilateral animals

September 23 2009

---



An international research team led by Brown University has determined that the flatworm *Acoelomorpha* belongs as a sister clade to other bilateral animals. The finding means the worm is a product of the deepest split within the bilateral animals, the first evolutionary divergence within the group. Credit: Eric Rottinger/Kahikai.org

When it comes to understanding a critical junction in animal evolution, some short, simple flatworms have been a real thorn in scientists' sides. Specialists have jostled over the proper taxonomic placement of a group of worms called *Acoelomorpha*. This collection of worms, which comprises roughly 350 species, is part of a much larger group called bilateral animals, organisms that have symmetrical body forms, including

humans, insects and worms. The question about acoelomorpha, was: Where do they fit in?

To scientists, acoelomorpha, has been enigmatic, a "rogue animal," said Casey Dunn, an [evolutionary biologist](#) at Brown University. "It has been wandering throughout the animal tree of life."

The worm wanders no more. Through a laborious genetic sequencing analysis, Dunn and an international team of scientists have settled the long-standing debate and determined that acoelomorpha belongs as a sister clade to other bilateral animals. The finding is significant, Dunn said, because it shows the worm is a product of the deepest split within the bilateral animals, the first evolutionary divergence within the group. Because of that, scientists have gained a key insight into the most recent [common ancestor](#) to bilaterians, a species that remains unknown.

The worm is "as distant as an animal can be in bilateria and still be a bilaterian," said Dunn, assistant professor of biology. "So, now we have two perspectives to (find out about) this common ancestor, the acoelomorphs and all the other bilateral animals."

The results appear online this week in the [Proceedings of the Royal Society B](#).

The team, composed of 17 scientists from the United States, France Germany, Sweden, Spain and the United Kingdom, had two more interesting findings:

- The debate appears to be over for *Xenoturbella*, a type of marine worm whose ancestral affiliation had been tossed between worms and mollusks. The researchers reported their [genetic analysis](#) shows diminishing evidence for placing xenoturbella within

*Deuterostomia*, one of the major groups within the animal kingdom. Coincidentally, it also shows that xenoturbella may be a close relative to acoelomorpha.

- *Cycliophora*, a single species discovered in 1994 that lives on the bristles surrounding the mouth of the Norway lobster *Nephrops norvegicus*, has found a home with *Entoprocta* and *Ectoprocta*. The researchers base their findings on an analysis that reached further into the genetic makeup of cycliophora than previous studies had done.

The team used a genetic sequencing technique called expressed sequence tags to carry out the phylogenetic studies. The aim of this approach, discussed in a study led by Dunn that appeared in *Nature* last year, is to analyze a large number of genes from a large number of animals. For this paper, the researchers looked at 1,487 genes, a 10-fold increase in the number of genes analyzed in previous studies. In all, the researchers logged 2.25 million processor hours on a supercomputer in California to obtain the results. Dunn called the effort the most computationally intensive phylogenetic analysis to date.

Source: Brown University ([news](#) : [web](#))

Citation: Research team finds first evolutionary branching for bilateral animals (2009, September 23) retrieved 1 May 2024 from <https://phys.org/news/2009-09-team-evolutionary-bilateral-animals.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.