

Researchers identify dominant chemical that attracts mosquitoes to humans

October 26 2009



A female mosquito of the Culicidae family (*Culiseta longiareolata*). Image: Wikipedia

Scientists at the University of California, Davis, have identified the dominant odor naturally produced in humans and birds that attracts the blood-feeding *Culex* mosquitoes, which transmit West Nile virus and other life-threatening diseases.

The groundbreaking research, published this week in the early online edition of the [Proceedings of the National Academy of Sciences](#), explains why mosquitoes shifted hosts from birds to humans and paves the way for key developments in mosquito and disease control.

Entomology professor Walter Leal and postdoctoral researcher Zain Syed found that nonanal (sounds like NAWN-uh-nawl) is the powerful semiochemical that triggers the mosquitoes' keen sense of smell, directing them toward a blood meal. A semiochemical is a chemical substance or mixture that carries a message.

"Nonanal is how they find us," Leal said. "The antennae of the *Culex quinquefasciatus* are highly developed to detect even extremely low concentrations of nonanal." Mosquitoes detect smells with the olfactory receptor neurons of their antennae.

Birds, the main hosts of mosquitoes, serve as the reservoir for the West Nile virus, Leal said. When infected mosquitoes take a blood meal, they transmit the virus to their hosts, which include birds, humans, horses, dogs, cats, bats, chipmunks, skunks, squirrels and domestic rabbits. Since 1999, the U.S. Centers for Disease Control and Prevention have recorded 29,397 human cases and 1,147 fatalities in the United States alone.

The UC Davis researchers tested hundreds of naturally occurring compounds emitted by people and birds. They collected chemical odors from 16 adult human subjects, representing multiple races and ethnic groups.

"We then determined the specificity and sensitivity of the olfactory receptor neurons to the isolated compounds on the antennae of the [mosquitoes](#)," Syed said.

Leal and Syed found that nonanal acts synergistically with [carbon dioxide](#), a known mosquito attractant. "We baited mosquito traps with a combination of nonanal and carbon dioxide and we were drawing in as many as 2,000 a night in Yolo County, near Davis," Syed said. "Nonanal, in combination with carbon dioxide, increased trap captures by more

than 50 percent, compared to traps baited with carbon dioxide alone."

Source: University of California - Davis

Citation: Researchers identify dominant chemical that attracts mosquitoes to humans (2009, October 26) retrieved 18 April 2024 from <https://phys.org/news/2009-10-dominant-chemical-mosquitoes-humans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.