

Fruit juices contain more vitamin C than their labels indicate

October 5 2009



Of the 17 samples analyzed, the one with the highest vitamin C content was apple juice. Credit: Leonid Mamchenkow

A team of pharmacists from the University of Santiago de Compostela (USC) has established that the levels of vitamin C in many fruit juices and soft drinks are far higher than those indicated on their labels by the manufacturers. This finding has been possible owing to a new technique developed by the researchers to determine the content of vitamin C in these kinds of drinks.

Ascorbic acid or vitamin C is a [natural antioxidant](#) in fruits and vegetables, but the European Commission permits its use as an additive in juices, jams, dairy products and other foods. The involvement of this

substance in the immune response and other biochemical processes such as the formation of collagen and the absorption of iron is well-known. However, high levels of [ascorbic acid](#) can cause diarrhoea and gastrointestinal problems, as a result of which scientists are attempting to determine the content of vitamin C in foods with greater and greater accuracy.

Now, a group of researchers from the Faculty of Pharmacy of the USC has developed a new chromatographic technique (these are used to separate and identify chemical elements) aimed at accurately measuring the ascorbic acid in fruit juices and soft drinks. By applying this method, they have found that the amounts of vitamin C stipulated on the labels of many drinks are not real. In a sample of 17 fruit juices, soft drinks and isotonic drinks, only two correspond to what is indicated on the bottle.

Ana Rodríguez Bernaldo de Quirós is a member of the team which has developed the new technique, whose details have recently been published in the *Food Chemistry* magazine. "The other drinks contain much higher levels than those specified by the manufacturer because, as has already been indicated in a previous study, the label probably only shows the amount of added ascorbic acid, without taking into account the fruit's natural vitamin C content", she explained to SINC.

Bernaldo de Quirós highlights the greater resolution and sensitivity of the method, by means of which it is possible to detect up to 0.01 milligrams of vitamin C per litre, "thanks to the use of new column chromatography, based on spherical particles of ultra pure silica 3 microns in size".

"Another advantage of the method is its simplicity and speed, as the total time taken to carry out the analyses is no more than six minutes", the researcher remarked.

With the new technique, the valuation of the ascorbic acid in the drinks has revealed some curious data. Of the 17 samples analyzed, the one with the highest vitamin C content was an apple juice (840 mg/l), more than the orange juices (352-739 mg/l). The results for the pineapple and grape juices were 702 mg/l and between 30.2 and 261 mg/l for the soft drinks (orange, lemon and apple).

The researchers also evaluated how the [vitamin C](#) content of the orange juices and tea drinks varies while they are on the shelves in the temperature conditions specified by the manufacturer. After six days, the former barely lose 8% of their ascorbic acid while, in the tea drinks, this substance falls by 54% at 4°C and practically disappears at room temperature.

More information: A. Rodríguez-Bernaldo de Quirós, M. Fernández-Arias, J. López-Hernández. "A screening method for the determination of ascorbic acid in fruit juices and [soft drinks](#)". *Food Chemistry* 116 (2): 509-512, 2009.

Source: FECYT - Spanish Foundation for Science and Technology

Citation: Fruit juices contain more vitamin C than their labels indicate (2009, October 5) retrieved 3 May 2024 from <https://phys.org/news/2009-10-fruit-juices-vitamin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.