

# Ecologists sound out new solution for monitoring cryptic species

November 27 2009

---

Ecologists have at last worked out a way of using recordings of birdsong to accurately measure the size of bird populations. This is the first time sound recordings from a microphone array have been translated into accurate estimates of bird species' populations. Because the new technique, reported in the British Ecological Society's *Journal of Applied Ecology*, will also work with whale song, it could lead to a major advance in our ability to monitor whale and dolphin numbers.

Developed by Deanna Dawson of the US Geological Survey and Murray Efford of the University of Otago, New Zealand, the technique is an innovative combination of sound recording with spatially explicit capture-recapture (SECR), a new version of one of ecologists' oldest tools for monitoring animal populations.

Birds communicate by singing or calling, and biologists have long counted these cues to get an index of bird abundance. But it is much harder to work out the actual density of a bird population because existing methods need observers to measure either the distance to each bird, or whether they are within a set distance from the observer. This is straightforward if birds are seen, but difficult when birds are heard but not seen.

According to Dawson: "We devised a way to estimate population density of birds or other animals that vocalise by combining sound information from several microphones. A sound spreading through a forest or other habitat leaves a 'footprint'. The size of the footprint depends on how

quickly the sound attenuates. Mathematically, there is a unique combination of population density and attenuation rate that best matches the number and 'size' of the recorded sounds. We used computer methods to find the best match, and thereby estimate density."

Dawson and Efford developed the method by recording the ovenbird - a warbler more often heard than seen - in deciduous forest at the Patuxent Research Refuge near Laurel, Maryland, USA. They rigged up four microphones close to the ground in a square with 21 metre-long sides. Over five days, they moved the microphones to 75 different points across their study area and recorded ovenbirds singing.

They chose the ovenbird as the species from which to develop the method because of its concise, distinctive song and because the males sing from the lower layers of the forest.

The new acoustic technique gives a more accurate estimate of bird numbers than using nets to capture birds, which can be stressful for the [birds](#) as well as time consuming for the researchers.

As well as helping assess populations of cryptic bird species such as the ovenbird, the new technique might be applied to measuring hard-to-reach populations of marine mammals, such as whales and dolphins. Developing ways of estimating whale and dolphin numbers acoustically is seen as critical for understanding these species' populations.

Recording the sounds has other benefits, too. "Sound intensity and other characteristics can be measured from the spectrogram - the graph of the sounds - to improve density estimates. Archiving the sounds also makes it possible to re-examine them, or to extract additional information as analytical methods evolve," says Dawson.

**More information:** Deanna K. Dawson and Murray G. Efford (2009).

Bird population density estimated from acoustic signals, [Journal of Applied Ecology](#), doi: [10.1111/j.1365-2664.2009.01731.x](https://doi.org/10.1111/j.1365-2664.2009.01731.x), is published online on 27 November 2009.

Source: Wiley ([news](#) : [web](#))

Citation: Ecologists sound out new solution for monitoring cryptic species (2009, November 27)  
retrieved 19 April 2024 from  
<https://phys.org/news/2009-11-ecologists-solution-cryptic-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.