

'Ultra-primitive' particles found in comet dust

November 2 2009



Artist concept of Stardust.

Dust samples collected by high-flying aircraft in the upper atmosphere have yielded an unexpectedly rich trove of relicts from the ancient cosmos, report scientists from the Carnegie Institution. The stratospheric dust includes minute grains that likely formed inside stars that lived and died long before the birth of our sun, as well as material from molecular clouds in interstellar space. This "ultra-primitive" material likely wafted into the atmosphere after the Earth passed through the trail of an Earthcrossing comet in 2003, giving scientists a rare opportunity to study cometary dust in the laboratory.



At high altitudes, most dust in the atmosphere comes from space, rather than the Earth's surface. Thousands of tons of interplanetary dust particles (IDPs) enter the atmosphere each year. "We've known that many IDPs come from comets, but we've never been able to definitively tie a single IDP to a particular comet," says study coauthor Larry Nittler, of Carnegie's Department of <u>Terrestrial Magnetism</u>. "The only known cometary samples we've studied in the laboratory are those that were returned from comet 81P/Wild 2 by the Stardust mission." The Stardust mission used a NASA-launched spacecraft to collect samples of comet dust, returning to Earth in 2006.

Comets are thought to be repositories of primitive, unaltered matter left over from the formation of the <u>solar system</u>. Material held for eons in cometary ice has largely escaped the heating and chemical processing that has affected other bodies, such as the planets. However, the Wild 2 dust returned by the Stardust mission included more altered material than expected, indicating that not all cometary material is highly primitive.

The IDPs used in the current study were collected by NASA aircraft in April 2003, after the Earth passed through the dust trail of comet Gregg-Skjellerup. The research team, which included Carnegie scientists Nittler, Henner Busemann (now at the University of Manchester, U.K.), Ann Nguyen, George Cody, and seven other colleagues, analyzed a subsample of the dust to determine the chemical, isotopic and microstructural composition of its grains. The results are reported online in *Earth and Planetary Science Letters*.





Scanning electron images of two dust particles E1 (panel A) and G4 (B) and secondary ion mass spectrometry isotopic ratio maps (C-D). Oxygen isotope maps of particles E1 (C) and G4 (D) show four and seven isotopically anomalous regions, indicated by circles, which have been identified as presolar grains. The scale bars are 2 microns.

"What we found is that they are very different from typical IDPs" says Nittler. "They are more primitive, with higher abundances of material whose origin predates the formation of the solar system." The distinctiveness of the particles, plus the timing of their collection after the Earth's passing through the comet trail, point to their source being the Gregg-Sk jellerup comet.

"This is exciting because it allows us to compare on a microscopic scale in the laboratory dust particles from different comets," says Nittler. "We can use them as tracers for different processes that occurred in the solar system four-and-a-half billion years ago."

The biggest surprise for the researchers was the abundance of so-called presolar grains in the dust sample. Presolar grains are tiny dust particles



that formed in previous generations of stars and in supernova explosions before the formation of the solar system. Afterwards, they were trapped in our solar system as it was forming and are found today in meteorites and in IDPs. Presolar grains are identified by having extremely unusual isotopic compositions compared to anything else in the solar system. But presolar grains are generally extremely rare, with abundances of just a few parts per million in even the most primitive meteorites, and a few hundred parts per million in IDPs. "In the IDPs associated with comet Gregg-Skjellerup they are up to the percent level," says Nittler. "This is tens of times higher abundances than we see in other primitive materials."

Also surprising is the comparison with the samples from Wild 2 collected by the Stardust mission. "Our samples seem to be much more primitive, much less processed, than the samples from Wild 2," says Nittler, "which might indicate that there is a huge diversity in the degree of processing of materials in different comets."

<u>More information:</u> Busemann, H., et al., Ultra-primitive interplanetary <u>dust particles</u> from the <u>comet</u> 26P/Grigg-Skjellerup dust stream collection, *Earth Planet. Sci. Lett.* (2009), <u>doi:10.1016/j.epsl.2009.09.007</u>

Source: Carnegie Institution

Citation: 'Ultra-primitive' particles found in comet dust (2009, November 2) retrieved 2 May 2024 from <u>https://phys.org/news/2009-11-ultra-primitive-particles-comet.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.