

# Brightness variations of sun-like stars: The mystery deepens

December 7 2009

---



Born from clouds of gas and dust, stars like our Sun spend most of their lifetime slowly burning their primary nuclear fuel, hydrogen, into the heavier element helium. After leading this bright and shiny life for several billion years, their fuel is almost exhausted and they start swelling, pushing the outer layers away from what has turned into a small and very hot core. These "middle-aged" stars become enormous, hence cool and red -- red giants. All red giants exhibit a slow oscillation in brightness due their rhythmic "breathing" in and out, and one third of them are also affected by additional, slower and mysterious changes in their luminosity. After this rapid and tumultuous phase of their later life, these stars do not end in dramatic explosions, but die peacefully as planetary nebulae, blowing out everything but a tiny remnant, known as white dwarf. Credit: ESO/S. Steinhöfel

(PhysOrg.com) -- An extensive study made with ESO's Very Large Telescope deepens a long-standing mystery in the study of stars similar

to the Sun. Unusual year-long variations in the brightness of about one third of all Sun-like stars during the latter stages of their lives still remain unexplained. Over the past few decades, astronomers have offered many possible explanations, but the new, painstaking observations contradict them all and only deepen the mystery. The search for a suitable interpretation is on.

"Astronomers are left in the dark, and for once, we do not enjoy it," says Christine Nicholls from Mount Stromlo Observatory, Australia, lead author of a paper reporting the study. "We have obtained the most comprehensive set of observations to date for this class of Sun-like stars, and they clearly show that all the possible explanations for their unusual behaviour just fail."

The mystery investigated by the team dates back to the 1930s and affects about a third of Sun-like stars in our Milky Way and other [galaxies](#). All stars with masses similar to our Sun become, towards the end of their lives, red, cool and extremely large, just before retiring as white dwarfs. Also known as [red giants](#), these elderly stars exhibit very strong periodic variations in their luminosity over timescales up to a couple of years.

"Such variations are thought to be caused by what we call 'stellar pulsations'," says Nicholls. "Roughly speaking, the [giant star](#) swells and shrinks, becoming brighter and dimmer in a regular pattern. However, one third of these stars show an unexplained additional periodic variation, on even longer timescales — up to five years."

In order to find out the origin of this secondary feature, the astronomers monitored 58 stars in our galactic neighbour, the [Large Magellanic Cloud](#), over two and a half years. They acquired [spectra](#) using the high resolution FLAMES/GIRAFFE [spectrograph](#) on ESO's Very Large Telescope and combined them with images from other telescopes, achieving an impressive collection of the properties of these variable

stars.

Outstanding sets of data like the one collected by Nicholls and her colleagues often offer guidance on how to solve a cosmic puzzle by narrowing down the plethora of possible explanations proposed by the theoreticians. In this case, however, the observations are incompatible with all the previously conceived models and re-open an issue that has been thoroughly debated. Thanks to this study, astronomers are now aware of their own "ignorance" — a genuine driver of the knowledge-seeking process, as the ancient Greek philosopher Socrates is said to have taught.

"The newly gathered data show that pulsations are an extremely unlikely explanation for the additional variation," says team leader Peter Wood. "Another possible mechanism for producing [luminosity](#) variations in a star is to have the star itself move in a binary system. However, our observations are strongly incompatible with this hypothesis too."

The team found from further analysis that whatever the cause of these unexplained variations is, it also causes the giant [stars](#) to eject mass either in clumps or as an expanding disc. "A Sherlock Holmes is needed to solve this very frustrating mystery," concludes Nicholls.

More information: This research was presented in two papers: one appeared in the November issue of the *Monthly Notices of the Royal Astronomical Society* ("Long Secondary Periods in Variable Red Giants", by C. P. Nicholls et al.), and the other has just been published in the *Astrophysical Journal* ("Evidence for mass ejection associated with long secondary periods in red giants", by P. R. Wood and C. P. Nicholls). Research papers: [arxiv.org/abs/0907.2975](http://arxiv.org/abs/0907.2975) and [arxiv.org/abs/0910.4418](http://arxiv.org/abs/0910.4418)

Source: ESO ([news](#) : [web](#))

Citation: Brightness variations of sun-like stars: The mystery deepens (2009, December 7)  
retrieved 1 May 2024 from

<https://phys.org/news/2009-12-brightness-variations-sun-like-stars-mystery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.