

Farms, Fertilizers and Greenhouse Gas Emissions

December 10 2009, By Ann Perry



Research by ARS soil scientist Rod Venterea on the release of nitrous oxide and other greenhouse gases suggests farmers using reduced tillage can minimize nitrous oxide emissions by putting fertilizers below the upper 2 to 3 inches of soil.

(PhysOrg.com) -- Agricultural Research Service (ARS) scientists are front and center in finding out how farming affects emissions of the green house nitrous oxide (N₂O).

Experts already know that N₂O emissions rise as applications of nitrogen-based fertilizers increase. Microbiologist Tim Parkin, who works at the ARS National Laboratory for Agriculture and the

Environment in Ames, Iowa, is part of a team that is studying how different soils and different fertilizers affect N₂O emissions.

The researchers assessed the variation in the emissions of N₂O, carbon dioxide and methane from two different soil types—a sandy loam mix and a clay soil. The two fertilizers used in the study were urea-ammonium nitrate (UAN) and a liquid swine manure slurry.

They found that overall N₂O emission levels were highest from soils amended with swine manure slurry. High levels of N₂O emissions were measured from sandy loam soils amended either with UAN or slurry. But on the clay soils, only those amended with slurry—and not with UAN—had elevated N₂O emissions.

Soil scientist Rod Venterea, who works at the ARS Soil and Water Management Research Unit in St. Paul, Minn., is also studying N₂O emission dynamics. He found that the amount of N₂O emitted from fields fertilized with anhydrous ammonia was on average twice as high as emissions from fields fertilized with urea. The higher emissions from anhydrous ammonia were likely derived from the conversion of ammonia to nitrate.

His findings also suggest that farmers using reduced tillage can minimize N₂O emissions by placing fertilizers below the upper 2 to 3 inches of soil. This is because in a reduced tillage system, the microorganisms that support N₂O emissions are concentrated in the topmost [soil](#) layer.

Results from Parkin's research were published in the Journal of Environmental Quality in 2008. Venterea's work was published in Global Change Biology in 2007 and the Journal of Environmental Quality in 2005 and 2008.

[Read more](#) about this research in the November/December 2009 issue of

Agricultural Research magazine.

Provided by USDA [Agricultural Research](#) Service

Citation: Farms, Fertilizers and Greenhouse Gas Emissions (2009, December 10) retrieved 2 May 2024 from <https://phys.org/news/2009-12-farms-fertilizers-greenhouse-gas-emissions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.