

First Direct Imaging of a Young Binary System

December 15 2009





Figure 1: Observed and simulated images of the young binary star SR24 (distance: 520 light years). A. Infrared image of SR24 and associated disks. The regions gravitationally bound to each star (Roche Lobes) are indicated by dotted lines (for the inner regions) and dashed lines (for the outer regions). (©The graduate University for Advanced Studies & the National Astronomical Observatory of Japan) B. Snapshot of the buildup of materials onto the binary system SR24 based on two-dimensional numerical simulations. The colors designate the distribution of surface densities and the arrows, the distribution of velocity. (© Chiba University)

(PhysOrg.com) -- A team of astronomers from The Graduate University for Advanced Studies, the National Astronomical Observatory of Japan, and other universities have captured the first direct image of a young

binary star system. Using the Coronagraphic Imager with Adaptive Optics (CIAO) mounted on the Subaru Telescope, the team observed the young binary star SR24, which is located in the constellation Ophiuchus, 520 light years away.


Our understanding of the creation and development of [stars](#) and [planets](#) has advanced substantially in the past two decades. Research has confirmed that new stars are often surrounded by disks of dense gas (“protoplanetary disks”) from which planets are believed to form; the radii of these disks can extend distances up to several hundreds times farther than the distance between the Sun and the Earth. Scientists have intensively studied the structure of protoplanetary disks at various electromagnetic wavelengths, and our knowledge of the mechanisms of single star formation has increased considerably. Nevertheless, most stars form in binary or multiple (more than two stars) systems, and many unanswered questions remain about how they are born and develop (see Figure 2 for a numerical simulation of binary formation and Figure 3 and the movie for the birth of a binary system). Consequently, studies of protoplanetary disks in binary and multiple systems are essential for making generalizations about the processes of star and [planet formation](#).



Three-dimensional numerical simulation image of young binary star system. (©

In a binary system, two stars orbit around a common center of mass. The brighter or more massive star is referred to as the primary star and the fainter or less massive one is called the secondary star. A disk of gas and dust surrounds the primary star (“circumprimary disk”); another disk encircles the secondary star (circumsecondary disk). A third disk (“circumbinary disk”) may ring the primary and secondary stars and supply mass to the stars’ disks through a stream of gas. Spiral arms of gas and dust may extend from the stars’ disks to feed them with material from the circumbinary disk, thus contributing to the development of their circumstellar disks. However, such disks and [spiral arms](#) in binary or multiple systems have rarely been directly imaged or resolved—until now.

The July 2006 observations of the young binary star SR24 by astronomers Dr. Satoshi Mayama (The Graduate University for Advanced Studies or SOKENDAI), Dr. Motohide Tamura (National Astronomical Observatory of Japan or NAOJ and SOKENDAI), Dr. Masahiko Hayashi (NAOJ and SOKENDAI) and colleagues have corrected this deficit and provided data that can be used to test theories about binary system formation. The team captured a high-resolution, near-infrared image of the circumprimary and circumsecondary disks of SR24, the first such image of twin protoplanetary disks around a young stellar object (Figure 1A, Figure 4). They also discovered a long spiral arm extending out from the circumprimary disk, another first in observations of young stars. The image shows a bridge of gas connecting the two disks. The observations provide the first data that can be used to test theoretical models of mass accretion in binary systems.



Left panel: Wide-field three-color composite image of the environment around the constellation Ophiuchus, where SR24 is located. (Obtained with the 1.4 m telescope IRSF [Infrared Survey Facility] at the South African Astronomical Observatory [SAAO]) (© The Graduate University for Advanced Studies) Right panel: Image of the young binary system SR24. Obtained with the 8.2m Subaru Telescope. (© The Graduate University for Advanced Studies)




Diagram of the birth of a binary (© Hosei University) The team observed the young binary SR24 in its formative stage and detected three disks for the first time. These are portrayed in the figure. Arrows denote the velocity distribution of materials in this figure. In a binary system, the brighter star is called the primary star and the fainter star, the secondary star. While a single star has only one circumstellar disk, a binary has three types of disks. In a binary system, both the primary and secondary stars orbit each other and have circumprimary and circumsecondary disks respectively. The entire system can be surrounded by a circumbinary disk. Numerical simulation leads to the assumption that there is a

stage when a young binary is associated with all three disks.

Astronomers Dr. Tomoyuki Hanawa (Chiba University) and Dr. Tomoaki Matsumoto (Hosei University) analyzed the young binary star and collected information about its attributes. Using a supercomputer, they performed two-dimensional numerical simulations of accretion from a circumbinary disk and were able to generate stellar structures (Figure 1B) that shared common features with the observed image. The agreement between the observations and the simulation suggests that fresh material streams along the spiral arm, replenishing the circumprimary disk with a reservoir of gas contained in a circumbinary disk. As reported in this article, the 0.1 arcsecond image shows the supporting structures of twin protoplanetary disks, which are fed with planet-building materials. Such an image cannot be reproduced by spectroscopic observations or studies of models of spectral energy distribution. The simulations also confirm that the bridge between the twin disks corresponds to gas flow and a shock wave caused by the collision of gas rotating around the primary and secondary stars. The comparison of the simulations with the observation demonstrates a mechanism for how protoplanetary disks interact in young binary systems.

These findings are a testament to the significance of this first direct observation of a binary system that is in the process of forming. They lead to a better understanding of the process of star and planet formation in a binary system by clarifying the role of supporting structures in maintaining the binary system.

More information: These findings were published in the November 19, 2009 edition of Science Express in an article entitled “Direct Imaging of Bridged Twin Protoplanetary Disks in a Young Multiple

Star” by S. Mayama et al. and will be subsequently published in Science.

Provided by Subaru Telescope

Citation: First Direct Imaging of a Young Binary System (2009, December 15) retrieved 25 April 2024 from <https://phys.org/news/2009-12-imaging-young-binary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.