

Levitating magnet may yield new approach to clean energy

January 24 2010, by David L. Chandler



The Levitated Dipole Experiment (LDX) reactor is housed inside a 16-foot-diameter steel structure in a building on the MIT campus that also houses MIT's other fusion reactor, a tokamak called Alcator C-mod.

(PhysOrg.com) -- A new experiment that reproduces the magnetic fields of the Earth and other planets has yielded its first significant results. The findings confirm that its unique approach has some potential to be developed as a new way of creating a power-producing plant based on nuclear fusion -- the process that generates the sun's prodigious output of energy.

Fusion has been a cherished goal of physicists and energy researchers for more than 50 years. That's because it offers the possibility of nearly endless supplies of energy with no carbon emissions and far less radioactive waste than that produced by today's nuclear plants, which are

based on fission, the splitting of atoms (the opposite of fusion, which involves fusing two atoms together). But developing a [fusion reactor](#) that produces a net output of energy has proved to be more challenging than initially thought.

The new results come from an experimental device on the MIT campus, inspired by observations from space made by satellites. Called the Levitated Dipole Experiment, or LDX, a joint project of MIT and Columbia University, it uses a half-ton donut-shaped magnet about the size and shape of a large truck tire, made of superconducting wire coiled inside a stainless steel vessel. This magnet is suspended by a powerful [electromagnetic field](#), and is used to control the motion of the 10-million-degree-hot electrically charged gas, or plasma, contained within its 16-foot-diameter outer chamber.

The results, published this week in the journal [Nature Physics](#), confirm the counter-intuitive prediction that inside the device's magnetic chamber, random turbulence causes the plasma to become more densely concentrated — a crucial step to getting atoms to fuse together — instead of becoming more spread out, as usually happens with turbulence. This "turbulent pinching" of the plasma has been observed in the way plasmas in space interact with the Earth's and Jupiter's magnetic fields, but has never before been recreated in the laboratory.

Most experiments in fusion around the world use one of two methods: tokamaks, which use a collection of coiled magnets surrounding a donut-shaped chamber to confine the plasma, or inertial fusion, using high-powered lasers to blast a tiny pellet of fuel at the device's center. But LDX takes a different approach. "It's the first experiment of its kind," says MIT senior scientist Jay Kesner, MIT's physics research group leader for LDX, who co-directs the project with Michael E. Mauel, professor of applied physics at Columbia University's Fu Foundation School of Engineering and Applied Science.

The results of the experiment show that this approach "could produce an alternative path to fusion," Kesner says, though more research will be needed to determine whether it would be practical. For example, though the researchers have measured the plasma's high density, new equipment still needs to be installed to measure its temperature, and ultimately a much larger version would have to be built and tested.

Kesner cautions that the kind of fuel cycle planned for other types of fusion reactors such as tokamaks, which use a mixture of two forms of "heavy" hydrogen called deuterium and tritium, should be easier to achieve and will likely be the first to go into operation. The deuterium-deuterium fusion planned for devices based on the LDX design, if they ever become practical, would likely make this "a second-generation approach," he says.

When operating, the huge LDX magnet is supported by the [magnetic field](#) from an electromagnet overhead, which is controlled continuously by a computer based on precision monitoring of its position using eight laser beams and detectors. The position of the half-ton magnet, which carries a current of one million amperes (compared to a typical home's total capacity of 200 amperes) can be maintained this way to within half a millimeter. A cone-shaped support with springs is positioned under the magnet to catch it safely if anything goes wrong with the control system.

Levitation is crucial because the magnetic field used to confine the plasma would be disturbed by any objects in its way, such as any supports used to hold the magnet in place. In the experimental runs, they recreated the same conditions with and without the support system in place, and confirmed that the confinement of the plasma was dramatically increased in the levitated mode, with the supports removed. With the magnet levitated, the central peak of plasma density developed within a few hundredths of a second, and closely resembled those observed in planetary magnetospheres (such as the magnetic fields

surrounding Earth and Jupiter).

Summarizing the difference between the two approaches, Kesner explains that in a tokamak, the hot plasma is confined inside a huge magnet, but in the LDX the magnet is inside the plasma. The whole concept, he says, was inspired by observations of planetary magnetospheres made by interplanetary spacecraft. In turn, he says, for planetary research the experiments in LDX can yield "a lot more subtle detail than you can get by launching satellites, and more cheaply."

The MIT and Columbia scientists say that if the turbulence-induced density enhancement exhibited by the LDX could be scaled up to larger devices, it might enable them to recreate the conditions necessary to sustain fusion reactions, and thus may point the way toward abundant and sustainable production of fusion energy.

"Fusion energy could provide a long-term solution to the planet's energy needs without contributing to global warming," says Columbia's Mauel.

The LDX project, led by Mauel and Kesner and funded by the U.S. Department of Energy, has been through more than 10 years of design, construction and testing, and produced its first experimental results in its levitated configuration last year, which are being reported in the analysis published this week. Dr. Darren Garnier of Columbia University, who directs LDX experimental operations, last month received the Rose Award for Excellence in Fusion Engineering for his work on LDX. A newly installed microwave interferometer array, developed by MIT graduate student Alex Boxer PhD '09, was used to make the precision measurements of the plasma concentrations that were used to observe the turbulent pinch.

"LDX is one of the most novel fusion plasma physics experiments underway today," says Stewart Prager, director of the Princeton Plasma

Physics Laboratory. Because of the unique geometry of the system, he says, "theoretical predictions indicate that the confinement of energy might be very favorable" for producing practical fusion power, but the theory needs to be confirmed in practice. "For these benefits to be realized, the somewhat bold theoretical predictions must be realized experimentally," he says.

Provided by Massachusetts Institute of Technology

Citation: Levitating magnet may yield new approach to clean energy (2010, January 24) retrieved 1 May 2024 from <https://phys.org/news/2010-01-levitating-magnet-yield-approach-energy.html>

| |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|