


A solid case of entanglement

January 11 2010



This is an SEM image of a typical Cooper pair splitter. The bar is 1 micrometer. A central superconducting electrode (blue) is connected to two quantum dots engineered in the same single wall carbon nanotube (in purple). Entangled electrons inside the superconductor can be coaxed to move in opposite directions in the nanotube, ending up at separate quantum dots, while remaining entangled. Credit: L.G. Herrmann, F. Portier, P. Roche, A. Levy Yeyati, T. Kontos, and C. Strunk

Physicists have finally managed to demonstrate quantum entanglement of spatially separated electrons in solid state circuitry.

For the first time, physicists have convincingly demonstrated that physically separated particles in solid-state devices can be quantum-mechanically entangled. The achievement is analogous to the [quantum entanglement](#) of light, except that it involves particles in circuitry instead of photons in optical systems. Both optical and solid-state entanglement offer potential routes to [quantum computing](#) and secure communications, but solid-state versions may ultimately be easier to

incorporate into electronic devices.

The experiment is reported in an upcoming issue of *Physical Review Letters* and highlighted with a Viewpoint in the January 11 issue of *Physics*.

In optical entanglement experiments, a pair of entangled photons may be separated via a beam splitter. Despite their physical separation, the entangled photons continue to act as a single quantum object. A team of physicists from France, Germany and Spain has now performed a solid-state entanglement experiment that uses electrons in a superconductor in place of photons in an optical system.

As conventional [superconducting materials](#) are cooled, the electrons they conduct entangle to form what are known as Cooper pairs. In the new experiment, Cooper pairs flow through a superconducting bridge until they reach a [carbon nanotube](#) that acts as the electronic equivalent of a beam splitter. Occasionally, the electrons part ways and are directed to separate [quantum dots](#) -- but remain entangled. Although the quantum dots are only a micron or so apart, the distance is large enough to demonstrate entanglement comparable to that seen in optical systems.

In addition to the possibility of using entangled electrons in solid-state devices for computing and secure communications, the breakthrough opens a whole new vista on the study of quantum mechanically entangled systems in solid materials.

More information: Carbon Nanotubes as Cooper-Pair Beam Splitters, L. G. Herrmann, F. Portier, P. Roche, A. Levy Yeyati, T. Kontos, and C. Strunk, *Phys. Rev. Lett.* 104, 026801 (2010) - Published January 11, 2010, [Download PDF](#)

Provided by American Physical Society

Citation: A solid case of entanglement (2010, January 11) retrieved 10 April 2024 from <https://phys.org/news/2010-01-solid-case-entanglement.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.