

# How the butterflies got their spots

February 5 2010

---



(PhysOrg.com) -- How two butterfly species have evolved exactly the same striking wing colour and pattern has intrigued biologists since Darwin's day. Now, scientists at Cambridge have found 'hotspots' in the butterflies' genes that they believe will explain one of the most extraordinary examples of mimicry in the natural world.

Heliconius, or passion-vine butterflies, live in the Americas - from the southern United States to southern South America. Although they cannot interbreed, *H. melpomene* and *H. erato* have evolved to mimic one another perfectly.

These delicate butterflies have splashes of red and yellow on their black wings, signaling to birds that they contain toxins and are extremely

unpalatable. They mimic one another's colour and pattern to reinforce these warning signals.

Scientists have studied these butterflies since the 1860s as a classic case of evolution in action, but only now is modern sequencing technology unlocking the underlying genetics.

The Cambridge-led team of researchers from UK and US universities, which has been breeding the butterflies in Panama for the past decade, has been searching for the genes responsible for the butterflies' wing patterns and the answer to the question of whether the same genes in two different species are responsible for the mimicry.

According to Dr Chris Jiggins of the Department of Zoology at the University of Cambridge, one of the authors of the study: "The mimicry is remarkable. The two species that we study - erato and melpomene - are quite distantly related, yet you can't tell them apart until you get them in your hand. The similarity is incredible - even down to the spots on the body and the minute details of the wing pattern."

That the two species have evolved to look exactly the same is due to predation by birds. "The birds will try anything that looks different in the hope that it's good, so they learn that certain wing patterns are unpalatable and avoid them, but anything that deviates slightly from what they've experienced before is more likely to be attacked," he explains.

These butterflies have been studied since Darwin's day because they are such a striking example of adaptation. For years, scientists have pondered whether when different species evolve to look the same, they share a common genetic mechanism.

According to Jiggins: "It's interesting because it tells us how flexible

evolution is. If you get the same wing pattern evolving independently in different populations, do you expect the same genes to be involved?"

Because there are thousands of genes in the butterflies' genome, most scientists felt it was unlikely that the same genes should be involved. But the results of this study suggest that this is, in fact, the case.

The new results - published today in two parallel papers in the journal *PLoS Genetics* - show that the regions of the genome associated with the wing patterns are very small - akin to genetic "hotspots".

"This tells us something about the limitations on evolution, and how predictable it is. Our results imply that despite the many thousands of genes in the genome there are only one or two that are useful for changing this colour pattern. It seems like evolution might be concentrated in quite small regions of the genome - or hotspots - while the rest of it does not change very much," says Jiggins.

This is not the only unexpected element of the study. The team was also surprised that the obvious candidate [genes](#) - such as those involved in colour or wing pattern in other species - do not seem to be involved in the passion-vine butterflies' mimicry.

According to Jiggins: "We think it's more likely to be some novel method of cellular signaling, which is quite intriguing and could be important in many other insect species."

The next stage of the research is to look at other traits, such as behaviour, because the [butterflies](#) have preferences for particular colours and use wing patterns to select mates. "It seems the same regions of the genome control this behaviour as well as the wing pattern. We'd like to understand this," he says.

The results are published in *PLoS Genetics* on 5 February 2010.

Provided by University of Cambridge

Citation: How the butterflies got their spots (2010, February 5) retrieved 28 April 2024 from <https://phys.org/news/2010-02-butterflies.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--