

Will earlier springs throw nature out of step?

February 9 2010



The recent trend towards earlier UK springs and summers has been accelerating, according to a study published today (9 February 2010) in the scientific journal *Global Change Biology*.

The collaborative study, involving scientists from 12 UK research institutions, universities and conservation organisations, is the most comprehensive and rigorous assessment so far of long-term changes in the seasonal timing (phenology) of biological events across marine, freshwater and terrestrial environments in the UK.

Led by Dr Stephen Thackeray and Professor Sarah Wanless of the Centre for Ecology & Hydrology, the research gathers together more than 25,000 long-term phenology trends for 726 species of plants and animals. More than 80% of trends between 1976 and 2005 indicate

earlier seasonal events. The study considers a diverse array of organisms including plankton, plants, insects, amphibians, fish, birds and mammals. On average, the seasonal timing of reproduction and population growth has become earlier by more than 11 days over the whole period, but change has accelerated in recent decades.

The research shows that there are large differences between species in the rate at which seasonal events have shifted. Changes have been most rapid for many organisms at the bottom of food chains, such as plants and the animals that feed upon them. Predators have shown slower overall changes in the seasonal timing of their life cycle events. However, the seasonal timing of reproduction is often matched to the time of year when food supply increases, so that offspring receive enough food to survive. A key question is whether animals higher up the food chain will react to the faster rates of change in the plants and animals they feed upon, or whether they will fail to do so and become less successful at rearing their offspring.

Dr Thackeray said, "This is the first time that data have been analysed with enough consistency to allow a meaningful comparison of patterns of changing seasonal timing in the UK among such a diverse range of [plants](#) and animals."

Professor Wanless said, "It is important to realise that this analysis doesn't identify which predator-prey relationships are most at risk of disruption due to changes in timing. What it does do is highlight that the recorded changes need urgent investigation, particularly for species with high economic or conservation importance."

Co-author Richard Smithers of the Woodland Trust said, "Phenology is 'the canary in the cage'. The results of this new study make real our changing climate and its potential to have profound consequences for the complex web of life."

Provided by Centre for Ecology & Hydrology

Citation: Will earlier springs throw nature out of step? (2010, February 9) retrieved 2 May 2024 from <https://phys.org/news/2010-02-earlier-nature.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.