


# Single photons step into the slow light

February 22 2010

---



(PhysOrg.com) -- European experts in nanotechnology, optoelectronics and quantum physics have advanced the generation, detection and manipulation of single photons in semiconductors. Their discoveries bring practical single-photon and quantum applications significantly closer.

The ultimate in semiconductor miniaturisation is to use single photons and [electrons](#) to carry and manipulate information in the form of quantum bits or qubits. Researchers from some of Europe's leading universities, public research institutes and industry-based labs have combined forces under the aegis of the EU-funded QPHOTON project to create new semiconductor-based devices that reliably and efficiently

emit, detect and allow the manipulation of single photons.

“Current single-photon sources are not very useful in terms of efficiency and quality,” says Johann Reithmaier, the coordinator of QPHOTON.

“We set out to address this - to strongly improve the efficiency of producing single photons without any background of other photons.”

The suite of devices QPHOTON developed may enable multiple applications including low power, highly integrated photonic circuits; practical [quantum cryptography](#) and teleportation; and, eventually, new routes to realise ultra-powerful quantum computers.

## **Pillars, cavities and (quantum) dots**

One of the consortium’s key goals was to improve the assembly and control of semiconductor-based [quantum dots](#).

Quantum dots are nano-sized structures that can confine electrons in three dimensions. By manipulating the size, shape and composition of the dots, researchers can gain very fine control over their electronic and [optical properties](#), for example triggering them electronically to emit single photons.

Until now, most quantum dots have been grown through self-assembly, which leaves the dots scattered randomly in a solution or across a surface.

The QPHOTON researchers devised a way to grow quantum dots exactly where they wanted them by first etching minute holes in a semiconductor substrate. Still, getting the dots to have the precise properties the team wanted turned out to be a major challenge. “The primary layer was optically dead,” says Reithmaier, so we needed additional tricks to overcome this.”

The researchers developed a series of cleaning and deposition steps that allowed them to grow high-functioning dots where they wanted them, and, in the process, achieved a new record in terms of the density of quantum-dot spacing.

More importantly, however, the researchers were able to link those quantum dots to other nano-structures in order to enhance and exploit their properties. They found ways to combine the precisely spaced dots with minute vertical pillars and horizontally aligned microcavities, both of which emitted single photons far more efficiently and reliably than the previous state of the art.

The pillars emit light perpendicularly to the surface of the semiconductor. They can be used as robust sources of single photons, which could for example be coupled to optical fibres for long-distance transmission.

The horizontal microcavities are extremely promising as key parts of highly integrated circuits that would use photons rather than electrons to process information. The researchers also developed an innovative vertical photonic crystal assembly to extract single photons from these circuits with more than 80 percent efficiency.

Another major innovation was the development of “photonic wires” - precisely assembled linear structures that advanced the state of the art by a factor of two in terms of the efficiency of transmitting single photons. “Pushing the coupling efficiency to about 85 percent is a major breakthrough for single photon emitters,” says Reithmaier.

This advance is particularly promising for the secure distribution of quantum keys, which enable ultra-secure data transmission. Only by sending one photon at a time can information be transmitted without any risk of it being compromised.

## Slowing light to a crawl

For the past ten years, researchers worldwide have been avidly exploring the field of slow light. A variety of nano-structures have been found to have the ability to slow light dramatically. Engineers hope to use this effect to create memories, buffers and switches for high-powered, energy-efficient optical computers.

The QPHOTON researchers chalked up another first by demonstrating a strong slow light effect in a semiconductor quantum dot material.

Most previous research with slow light used media - gas for example - that would not be useful in actual computers. So showing that quantum dot arrays can slow light in a semiconductor is potentially very important for optical computing.

Reithmaier cautions that the ability of slow light to carry enough information to support optical data processing has not yet been demonstrated, and may turn out not to be possible. Still, QPHOTON's findings will be utilised in a new EU-funded project aimed at further exploring the potentials of slow light.

Reithmaier believes that the steps they've taken to control the emission, manipulation, and reception of single photons bring practical quantum computers significantly closer.

“The major challenge is to move quantum information from one location to another, and a major approach is single [photons](#),” he says. “For that you need full control of the photon, and that’s where we really pushed the technology.”

**More information:** QPHOTON project -- [www.ina.uni-kassel.de/qphoton](http://www.ina.uni-kassel.de/qphoton)

## Provided by ICT Results

Citation: Single photons step into the slow light (2010, February 22) retrieved 9 April 2024 from <https://phys.org/news/2010-02-photons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.