


Scientists Identify Specific Barley Tissues Infected by Scab

February 23 2010, By Stephanie Yao



By tracking where transformed *Fusarium graminearum* glows neon-green under a fluorescence microscope, ARS plant molecular geneticist Ron Skadsen has discovered that the fungus, which causes scab in barley, specifically attacks the protruding tip of the developing seed, the soft tissue connected with it and, to a lesser extent, the seed's outer hull.

(PhysOrg.com) -- Recent studies involving the fungus that causes the disease known as scab in barley have helped Agricultural Research Service (ARS) scientists identify the specific tissues the fungus infects.

The fungus *Fusarium graminearum* is the culprit behind [scab](#) in barley and wheat grain crops. Scab reduces yield by causing sterility and shrunken kernels, and contaminates the grain with mycotoxins.

Plant molecular geneticist Ron Skadsen at the ARS Cereal Crops Research Unit in Madison, Wis., infected barley seed spikes with

Fusarium transformed to contain a [green fluorescent protein](#) that makes the fungus glow neon-green when examined under a [fluorescence microscope](#). He found that Fusarium attacks the protruding seed tip of the developing seed, the soft tissue connected with it (just under the hull) and, to a lesser extent, the seed's outer hull.

Previous studies using detached barley seeds found that the fungus will liquefy the starchy interior of the seed within five days after inoculation. Skadsen found that, even 16 days after infection, Fusarium doesn't penetrate the starchy [endosperm](#) when the seed remains attached to the spike.

Knowing which parts of the barley plant Fusarium attacks gives insight into how the infection process works and where to target breeding and biotechnology strategies, according to Skadsen. Breeders can now focus on looking for traits that will prevent the [fungus](#)' early penetration.

These findings helped Skadsen and his research team to develop gene promoters that can be used to activate genes that defend against Fusarium in barley's susceptible tissues. Skadsen and research leader Cynthia Henson are now investigating the biochemistry of susceptible barley tissues through metabolic profiling during the first three days of infection.

[Read more](#) about this research in the February 2010 issue of *Agricultural Research* magazine.

Provided by USDA Agricultural Research Service

Citation: Scientists Identify Specific Barley Tissues Infected by Scab (2010, February 23) retrieved 16 July 2024 from <https://phys.org/news/2010-02-scientists-specific-barley-tissues-infected.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.