

F is for Fluoresence and Fluorine: New dyes for optical nanoscopy

March 26 2010

(PhysOrg.com) -- The imaging of living cells at the molecular level was barely a dream twenty years ago. Today, however, this dream is close to becoming reality.

In the Max Planck Institute for NanoBiophotonics in Göttingen, Stefan Hell (2009 recipient of the Otto Hahn Prize) has developed <u>fluorescence</u> <u>microscopy</u> methods for observing objects on the <u>nanoscale</u> and with his colleagues Vladimir Belov and Christian Eggeling a new series of photostable dyes that can be used as fluorescent markers has been realised, as reported in a cover story in *Chemistry -- A European Journal*.

Over the last two decades Stefan Hell and his group have revolutionized the art of microscopy beyond limits thought to have been unbreakable. Due to the wave properties (diffraction) of light, the resolution of an <u>optical microscope</u> is limited to object details of about 0.2 micrometers. The laws of physics appeared to prohibit imaging details beyond this limit. Stefan Hell saw beyond this limitation and about fifteen years ago his vision became concrete; he developed a method for observing objects at the nanometer scale by sequentially turning the fluorescence of nearby molecules off by stimulated emission, a technique known as STED nanoscopy.

The sensitivity of this technique depends on the brightness of the applied fluorescence markers and their photostability is also of great importance. The NanoBiophotonics group has succeeded in synthesizing a series of highly photostable and highly <u>fluorescent dyes</u>. These compounds emit



green and orange light and are based on <u>fluorine</u> derivatives of the wellknown Rhodamine dye. The use of these dyes in STED nanoscopy leads to images of high-quality with respect to brightness and signal-tobackground ratio; further the resolution over that of more traditional optical microscopes is significantly improved giving more detailed structural information.

These rhodamine-based fluorine derivatives are even more special because of their versatility. The compounds are available in hydrophilic and lipophilic forms, and with the inclusion of amino reactive groups, they can be easily attached to antibodies or other biomolecules in the course of standard labeling and immunostaining procedures. The group demonstrate that these new dyes are able to cross cellular membranes and reach the interior of living cells, which could lead to new labeling strategies for biological systems. All eyes are now on Göttingen to see just how far optical nanoscopy can go.

More information: Stefan Hell, New Fluorinated Rhodamines for Optical Microscopy and Nanoscopy, *Chemistry - A European Journal* 2010, 16, No. 15, 4477-4488, <u>dx.doi.org/10.1002/chem.200903272</u>

Provided by Wiley

Citation: F is for Fluoresence and Fluorine: New dyes for optical nanoscopy (2010, March 26) retrieved 2 May 2024 from https://phys.org/news/2010-03-fluoresence-fluorine-dyes-optical-nanoscopy.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.