

'Stick men' may be rendered obsolete in insect world

March 24 2010



Dr Mary Morgan-Richards with a female stick insect.

(PhysOrg.com) -- Male stick insects are becoming increasingly redundant, with new research showing some New Zealand female stick insects can reproduce as efficiently on their own as with a male mate.

Not only that, the females capable of reproducing without male help, always produce female [offspring](#).

The research, by Dr Mary Morgan-Richards of the Ecology Group, describes the distribution and [evolutionary relationships](#) of sexual and asexual populations of the stick insect.

The work was a collaboration with Dr Steve Trewick, also of the Ecology Group, and Dr Ian Stringer, from the Department of

Conservation in Wellington. The findings featured in a paper entitled Geographic parthenogenesis and the common tea-tree stick insect of New Zealand, published in the International journal **Molecular Ecology**.

The team conducted several experiments involving the reproduction of the species. Some populations have equal numbers of males and females that reproduce sexually, but others have unmated females that lay eggs that hatch and produce offspring identical to the mother, a process known as parthenogenesis.

The research also indicates that all of the southern parthenogenic populations seem to have the same [common ancestor](#). “That was unexpected,” Dr Morgan-Richards says. “It seems extraordinary when any single female is capable of reproducing parthenogenically.”

The team took females out of sexual populations and raised them on their own. Despite the lack of a male to mate with, the stick insects that usually reproduced sexually were capable of reproducing asexually.

“All of the parthenogenic populations of stick insect are to the south in New Zealand compared to the sexual populations more to the north,” Dr Morgan-Richards says. They don’t have a clear idea exactly of why that is but it seems to fit with the idea of range expansion—organisms moving to warmer places further north when the climate cools and expanding their population mid-range by going south when the climate warms.

The researchers found similar hatching success in mated and unmated females. Mated females produce equal numbers of male and female offspring, with most hatching within 9-16 weeks. In contrast, most of the offspring of unmated females were female, and the eggs took 21-23 weeks to hatch.

The difference in the development rate of the eggs “may have to do with the mechanism that they use to grow without sperm—but we don’t know”, Dr Morgan-Richards says. She suggests that competition between the two sexual forms could be influenced by an extended development rate in the south.

Females from sexual populations that had access to mates did not reproduce asexually, even though they were capable of doing so. Also, [females](#) from parthenogenic populations were able to reproduce sexually if they were given a male, but only about 10 per cent of their offspring were the result of sex. “It seemed that reverting to being sexual isn’t an easy step.”

The next phase of the research will try to determine the cause of such low rates of sexual reproduction from formerly parthenogenic populations.

Provided by Massey University

Citation: 'Stick men' may be rendered obsolete in insect world (2010, March 24) retrieved 27 April 2024 from <https://phys.org/news/2010-03-men-obsolete-insect-world.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
