


Answers to Probing Questions: Scientists pioneer new analysis that tells how uranium oxides stick together

March 16 2010



Reprinted with permission from Herman Cho, Wibe A. de Jong, and Chuck Z. Soderquist, J. Chem. Phys. 132, 084501. Copyright 2010, American Institute of Physics.

(PhysOrg.com) -- Combining experimental and theoretical methods, scientists at the Pacific Northwest National Laboratory gained a more complete picture of the electronic structure of a stable, soluble uranium ion and the bonds it forms. This study included an important first: the first nuclear magnetic resonance (NMR) spectroscopy data for solid UO_2^{2+} . Previous studies examined the ion in the liquid state. The new data provided insights into how the elements bond together to form

molecules and how the molecules combine to form clusters.

In addition, the team described tests that allow chemists to evaluate different theories on the bonds formed by this [uranium](#) ion, known as the uranyl ion or UO_2^{2+} . Selecting the most effective theories means scientists can focus experimental resources on the most promising ideas. Because of the significance of this research, it graced the February 28, 2010, cover of the [Journal of Chemical Physics](#).

This work advances the fundamental understanding of uranium's chemical interactions. Hence, this research directly contributes to predicting and manipulating uranium chemistry in the environment, in nuclear waste, and in biological systems. Intricate knowledge of toxins allows scientists and engineers to optimize strategies for environmental remediation and for storing nuclear waste, protecting human and ecosystem health.

To study the electronic structure of the uranyl ion, UO_2^{2+} , the researchers chose two solid salts. The first is an ammonium, carbonate, uranium combination, known by the formula $(\text{NH}_4)_4\text{UO}_2(\text{CO}_3)_3$. This salt was selected to shed light on hydrogen bonding to [oxygen atoms](#) in the uranyl ion. The second salt is rutherfordine, known by the formula UO_2CO_3 . It was selected as a control to exemplify a uranyl environment free of [hydrogen atoms](#).

The researchers conducted a series of NMR experiments, using a 300-MHz Tecmag Discovery NMR spectrometer, modified for analysis of radioactive samples. They carefully compared the NMR results to the results of theoretical calculations, specifically density functional theory calculations. The comparison revealed directions to a better understanding of bonding in uranium-containing molecules.

"This was a team effort with the experimentalists and theoreticians," said

Dr. Herman Cho, an experimental chemist at PNNL. "The theoreticians are getting ahead of experimentalists in calculating electronic interactions. Through this and other projects, we are trying to come up with experimental data to evaluate their models."

And, this project did just that. It results in a discriminating test for theoretical descriptions of covalent bonding involving heavy elements like uranium that contain high numbers of protons. This validates the approach used in this study and allows it to be extended to other molecules.

More information: Cho HM, WA De Jong, and CZ Soderquist. 2010. "Probing the Oxygen Environment in UO_2^{2+} by Solid-State O-17 NMR Spectroscopy and Relativistic Density Functional Calculations." *Journal of Chemical Physics* 132, 084501. [DOI:10.1063/1.3308499](https://doi.org/10.1063/1.3308499)

Provided by Pacific Northwest National Laboratory

Citation: Answers to Probing Questions: Scientists pioneer new analysis that tells how uranium oxides stick together (2010, March 16) retrieved 20 March 2023 from <https://phys.org/news/2010-03-probing-scientists-analysis-uranium-oxides.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.