

Southern Ocean winds open window to the deep sea

March 15 2010

Australian and US scientists have discovered how changes in winds blowing on the Southern Ocean drive variations in the depth of the surface layer of sea water responsible for regulating exchanges of heat and carbon dioxide between the ocean and the atmosphere.

The researchers' findings - published on-line today in *Nature Geoscience* - provide new insights into natural processes which have a major influence on the rate of <u>climate change</u>.

The surface-mixed layer is a crucial pathway between the atmosphere and the deeper layers of the ocean. Changes in the depth of the mixed layer can affect air-sea exchange, carbon and heat storage in the ocean, and the rate at which water sinks from the surface ocean into the deep ocean.

Changes in the mixed layer also affect <u>biological productivity</u>, by altering how much light and nutrients are available to support growth of <u>plankton</u> at the base of the food chain.

The paper's lead author, CSIRO Wealth from Oceans Flagship oceanographer Dr Jean-Baptiste Sallée, said the winds over the <u>Southern</u> <u>Ocean</u> had increased in strength and shifted closer to Antarctica in recent decades.

"The shift in winds is one of the strongest trends in southern hemisphere climate over the last 30 years," Dr Sallée said. "The key question is;



'How does the wind change affect the ocean?'

"Our knowledge of how the Southern Ocean changes in time is poor because of the lack of ship-based observations in this remote region. But we now have seven years of year-round observations from a fleet of profiling floats known as Argo, which allow us to see for the first time how the Southern Ocean changes with the seasons and from year-toyear."

The researchers, including Dr Steve Rintoul from the Antarctic Climate and Ecosystems CRC and CSIRO and Professor Kevin Speer from Florida State University, examined the relationship between changes in wind and changes in the surface-mixed layer.

"We found that the depth of the mixed layer was more sensitive than we expected to a <u>wind</u> pattern known as the Southern Annular Mode, the major mode of variability of the southern hemisphere atmosphere," Dr Sallée said. "Even more surprising was the fact that the response is very different in different regions."

When the winds strengthen and contract closer to Antarctica, the surfacemixed layer deepens in the eastern Indian and central Pacific oceans, and shallows in the western part of these basins. The reverse is seen when the winds weaken and migrate north.

The asymmetry can be explained by small deviations in the generally west-to-east winds and their effect on the heat exchange between ocean and atmosphere: when cold winds blow from the south, this causes heat loss from the ocean and deeper mixed layers.

"These changes in mixed layer depth affect how much light is available to support the growth of phytoplankton. We found that changes in the mixed layer depth driven by the winds are associated with changes in the



amount of phytoplankton biomass," Dr Sallée said.

Provided by CSIRO

Citation: Southern Ocean winds open window to the deep sea (2010, March 15) retrieved 2 May 2024 from <u>https://phys.org/news/2010-03-southern-ocean-window-deep-sea.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.