

Explained: Why many surveys of distant galaxies miss 90 percent of their targets

March 24 2010



This composite image of the GOODS-South field — the result of an extremely deep survey using two of the four giant 8.2-metre telescopes composing ESO's Very Large Telescope (VLT) and a unique custom-built filter — shows some of the faintest galaxies ever seen. It also allows astronomers to determine that 90% of galaxies whose light took 10 billion years to reach us have gone undiscovered. The image is based on data acquired with the FORS and HAWK-I instruments on the VLT. It shows in particular two varieties of light emitted by excited hydrogen atoms, known as Lyman-alpha and H-alpha. Credit: ESO/M. Hayes

(PhysOrg.com) -- Astronomers have long known that in many surveys of the very distant Universe, a large fraction of the total intrinsic light was not being observed. Now, thanks to an extremely deep survey using two



of the four giant 8.2-metre telescopes that make up ESO's Very Large Telescope (VLT) and a unique custom-built filter, astronomers have determined that a large fraction of galaxies whose light took 10 billion years to reach us have gone undiscovered. The survey also helped uncover some of the faintest galaxies ever found at this early stage of the Universe.

Astronomers frequently use the strong, characteristic "fingerprint" of light emitted by <u>hydrogen</u> known as the Lyman-alpha line, to probe the amount of stars formed in the very distant <u>Universe</u>. Yet there have long been suspicions that many distant <u>galaxies</u> go unnoticed in these surveys. A new VLT survey demonstrates for the first time that this is exactly what is happening. Most of the Lyman-alpha light is trapped within the galaxy that emits it, and 90% of galaxies do not show up in Lyman-alpha surveys.

"Astronomers always knew they were missing some fraction of the galaxies in Lyman-alpha surveys," explains Matthew Hayes, the lead author of the paper, published this week in *Nature*, "but for the first time we now have a measurement. The number of missed galaxies is substantial."

To figure out how much of the total luminosity was missed, Hayes and his team used the FORS camera at the VLT and a custom-built narrowband filter to measure this Lyman-alpha light, following the methodology of standard Lyman-alpha surveys. Then, using the new HAWK-I camera, attached to another VLT Unit <u>Telescope</u>, they surveyed the same area of space for light emitted at a different <u>wavelength</u>, also by glowing hydrogen, and known as the H-alpha line. They specifically looked at galaxies whose light has been travelling for 10 billion years (redshift 2.2), in a well-studied area of the sky, known as the GOODS-South field.



"This is the first time we have observed a patch of the sky so deeply in light coming from hydrogen at these two very specific wavelengths, and this proved crucial," says team member Göran Östlin. The survey was extremely deep, and uncovered some of the faintest galaxies known at this early epoch in the life of the Universe. The astronomers could thereby conclude that traditional surveys done using Lyman-alpha only see a tiny part of the total light that is produced, since most of the Lyman-alpha photons are destroyed by interaction with the interstellar clouds of gas and dust. This effect is dramatically more significant for Lyman-alpha than for H-alpha light. As a result, many galaxies, a proportion as high as 90%, go unseen by these surveys. "If there are ten galaxies seen, there could be a hundred there," Hayes says.

Different observational methods, targeting the light emitted at different wavelengths, will always lead to a view of the Universe that is only partially complete. The results of this survey issue a stark warning for cosmologists, as the strong Lyman-alpha signature becomes increasingly relied upon in examining the very first galaxies to form in the history of the Universe. "Now that we know how much light we've been missing, we can start to create far more accurate representations of the cosmos, understanding better how quickly stars have formed at different times in the life of the Universe," says co-author Miguel Mas-Hesse.

The breakthrough was made possible thanks to the unique camera used. HAWK-I, which saw first <u>light</u> in 2007, is a state-of-the-art instrument. "There are only a few other cameras with a wider field of view than HAWK-I, and they are on telescopes less than half the size of the VLT. So only VLT/HAWK-I, really, is capable of efficiently finding galaxies this faint at these distances," says team member Daniel Schaerer.

More information: This research was presented in a paper to appear in *Nature* ("Escape of about five per cent of Lyman-a photons from highredshift star-forming galaxies", by M. Hayes et al.).



Provided by ESO

Citation: Explained: Why many surveys of distant galaxies miss 90 percent of their targets (2010, March 24) retrieved 26 April 2024 from <u>https://phys.org/news/2010-03-surveys-distant-galaxies-percent.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.