

Hamlet fish sheds light on evolution of marine species

April 22 2010

To evolve or not to evolve? That is the question scientists at the University of East Anglia (UEA) are closer to answering following a groundbreaking new study into the colourful hamlet fish.

Little is known about the way [species](#) evolve in the vast marine environment. The new findings, published online today by the journal *Global Ecology and Biogeography*, significantly improve our understanding of speciation - the evolutionary process by which new species arise.

The hamlets are a group of colourful coral reef fish found throughout the Caribbean. Ten species of hamlet have been discovered and each can be easily recognized by its own distinct colour pattern. In some areas, as many as seven varieties can be found on a single reef. However, most hamlet species are only found at specific locations. The blue hamlet, for example, is found only in the Florida region.

It had previously been believed that these different species evolved because of geographical separation. For example, it was thought that falling sea levels in the past could have divided the original species. Then, when levels increased, the differently evolved species were thrown back together.

The new study by ecologists at UEA and Simon Fraser University in Canada found little evidence for this theory and instead suggests that hamlet colour varieties could have evolved regardless of any physical

separation.

Using thousands of underwater surveys made by SCUBA divers as part of the REEF volunteer survey project (www.reef.org), the researchers analysed distributions of the ten different hamlet species. They found that even widespread hamlet species are not found everywhere, and identified high density hotspots for each species.

Since different species hotspots overlap and many species have more than one hotspot, the results do not support the theory that hamlets originated independently when they were geographically separated in the past. The research also showed how ecological factors, such as competition for food or habitat, may influence how different hamlet species co-exist.

"Our findings suggest that ecology may better explain the evolution of hamlets than geographical separation," said lead author Dr Ben Holt of UEA's School of Biological Sciences.

"Many scientists believe hamlets are beginning to evolve into a new species and this latest discovery will shed light on this process."

More information: 'Signatures of speciation? Distribution and diversity of *Hypoplectrus* (Teleostei: Serranidae) colour morphotypes' by B Holt (UEA), I Cote (Simon Fraser University, Canada) and B Emerson (UEA) is published by Global Ecology and Biogeography on April 23.

Provided by University of East Anglia

Citation: Hamlet fish sheds light on evolution of marine species (2010, April 22) retrieved 9 April 2024 from <https://phys.org/news/2010-04-hamlet-fish-evolution-marine-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.