

## Magnetic 'ropes' tie down solar eruptions (w/ Video)

April 12 2010



The three images reveal gases trapped in the flux rope at different temperatures, from 1.5 million degrees Celsius in the image on the left through to 2.5 million degrees Celsius in the right hand image. These were made with data taken by the EIS telescope, an instrument built by a team led by UCL-MSSL and deployed on the Hinode spacecraft (a joint JAXA/UK/NASA mission). Credit: JAXA/ISAS/NASA/STFC

(PhysOrg.com) -- Using data from the Hinode spacecraft, a team of researchers from University College London (UCL) have revealed an immense magnetic structure that erupted to produce a dramatic solar eruption in late 2007. Solar scientist Dr Lucie Green will present their results at the RAS National Astronomy Meeting in Glasgow on Monday 12th April.

Over the last century, astronomers have become very aware of how just dynamic the Sun really is. One of the most dramatic manifestations of this is a coronal mass ejection (CME) where billions of tons of matter is thrown into space. If a CME reaches the Earth it creates inclement



'<u>space weather</u>' that can disrupt communications, power grids and the delicate systems on orbiting satellites. This potential damage means there is a keen interest in understanding exactly what triggers a CME outburst.

Now a team of researchers from University College London (UCL) have used data from the Hinode spacecraft to reveal new details of the formation of an immense magnetic structure that erupted to produce a CME on the 7th December 2007. Lead researcher Dr Lucie Green will present their results on Monday 12th April at the RAS National Astronomy Meeting in Glasgow.

The Sun's behaviour is shaped by the presence of magnetic fields that thread through the <u>solar atmosphere</u>. The magnetic fields may take on different shapes from uniform arches to coherent bundles of field lines known as 'flux ropes'. Understanding the exact structure of magnetic fields is a crucial part of the effort to determine how the fields evolve and the role they play in solar eruptions. In particular, flux ropes are thought to play a vital role in the CME process, having been frequently detected in interplanetary space as CMEs reach the vicinity of the Earth.

Dr. Green says, "Magnetic flux ropes have been observed in interplanetary space for many years now and they are widely invoked in theoretical descriptions of how CMEs are produced. We now need observations to confirm or reject the existence of flux ropes in the solar atmosphere before an eruption takes place to see whether our theories are correct."

The formation of the flux rope requires that significant energy is stored in the solar atmosphere. The rope is expected to remain stable whilst the solar <u>magnetic field</u> in the vicinity holds it down.

But at some point the structure becomes unstable and it erupts to produce a CME. Using data from the Hinode spacecraft Dr. Green has



shown that a flux rope formed in the solar atmosphere over the 2.5 days that preceded the December 2007 event. Evidence for the flux rope takes the form of S shaped structures which are clearly seen by one of the Hinode instruments, the UK-led Extreme-Ultraviolet Imaging Telescope.

The key point to understanding and predicting the formation of CMEs is to know when the flux rope becomes unstable. Combining the observations of the S shaped structure with information on how the magnetic field in the region evolves has enabled Dr.Green to work out when this happened. The work shows that over 30% of the magnetic field of the region had been transformed into the flux rope before it became unstable, three times what has been suggested in theory.

Dr Green sees a better understanding of magnetic flux ropes and their role in emissions from the Sun and other stars as one of the most pressing questions not just for solar physics but astronomy as a whole.

She comments, "Flux ropes are thought to play a vital role in the evolution of the magnetic field of the <u>Sun</u>. However, the physics of flux ropes is applied across the Universe. For example, a solar physics model of flux rope ejection was recently used to explain the jets driven by the accretion disks around the supermassive black holes found in the centre of galaxies."

## Provided by Royal Astronomical Society

Citation: Magnetic 'ropes' tie down solar eruptions (w/ Video) (2010, April 12) retrieved 5 May 2024 from <u>https://phys.org/news/2010-04-magnetic-ropes-solar-eruptions-video.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.