

# New mechanism for superconductivity discovered in iron-based superconductors

April 22 2010





Figure 1: Electronic standing waves in Fe(Se,Te) imaged by scanning tunnelling microscopy. Electronic standing waves are imaged as periodic horizontal and vertical streaks superposed on an inhomogeneous background (left column). Using a mathematical technique known as Fourier transformation, standing waves can be decomposed into components (right column). Bright spots in the Fourier maps, representing scatterings with particular characteristics, exhibit strong magnetic field dependence.

(PhysOrg.com) -- A research team at RIKEN, Japan's flagship research organisation has experimentally determined the mechanism underlying the formation of electron pairs in iron-based high-temperature

superconductors. The landmark finding, reported in the April 23rd issue of *Science*, establishes a key role for magnetism in superconductivity.

In classical theory, [superconductivity](#) occurs when two electrons are bound together to form a pair, known as a Cooper pair, by lattice vibrations. This pairing mechanism, however, has never been confirmed for high-temperature superconductors, whose transition temperatures well above the theoretical limit of about 40 K pose an enigma for condensed matter physics.

The iron-based superconductors investigated by the research team, first discovered in 2008 by Japanese researchers, offer the greatest chance of solving this enigma. With a maximum [transition temperature](#) of 55K, these superconductors are governed by an electron pairing mechanism that is different from earlier [superconductors](#) mediated by lattice vibrations, one based on two types of electrons with different momenta.



Magnetic-field induced change in the intensity of electronic standing waves evidencing the “ $s_{\pm}$ -wave” structure. When magnetic field is applied there appear two types of spots; one is enhanced by the field (blue) and the other is suppressed by the field (red). This behavior is evidence of the “ $s_{\pm}$ -wave” structure of Cooper pairs, strongly suggesting a magnetism-related pairing

mechanism.

To analyze this complex pairing mechanism, the researchers applied scanning tunnelling microscopy to electron pairing in Fe(Se, Te), the iron-based superconductor with the simplest crystal structure. Imaging electronic standing waves produced by scattering interference under a powerful 10-Tesla magnetic field, they found that Cooper pairs adopted a characteristic “ $s_{\pm}$ -wave” structure that is unique to a material with two types of electrons.

The discovery of  $s_{\pm}$ -wave structure breaks new ground by supporting a mechanism for electron pairing based not on lattice vibrations, as in other forms of superconductivity, but on magnetism. In providing a powerful constraint on [theoretical models](#), the finding thus marks a major advance toward unraveling the mystery of high-temperature superconductivity.

Provided by RIKEN

Citation: New mechanism for superconductivity discovered in iron-based superconductors (2010, April 22) retrieved 20 April 2024 from <https://phys.org/news/2010-04-mechanism-superconductivity-iron-based-superconductors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--