

# Triton's summer sky of methane and carbon monoxide

April 7 2010

---



This is an artist's impression of how Triton, Neptune's largest moon, might look from high above its surface. The distant Sun appears at the upper-left and the blue crescent of Neptune right of centre. Using the CRIRES instrument on ESO's Very Large Telescope, a team of astronomers has been able to see that the summer is in full swing in Triton's southern hemisphere. Credit: ESO/L. Calçada

According to the first ever infrared analysis of the atmosphere of Neptune's moon Triton, summer is in full swing in its southern hemisphere. The European observing team used ESO's Very Large Telescope and discovered carbon monoxide and made the first ground-based detection of methane in Triton's thin atmosphere. These observations revealed that the thin atmosphere varies seasonally, thickening when warmed.

"We have found real evidence that the [Sun](#) still makes its presence felt on Triton, even from so far away. This icy moon actually has seasons

just as we do on [Earth](#), but they change far more slowly," says Emmanuel Lellouch, the lead author of the paper reporting these results in *Astronomy & Astrophysics*.

On Triton, where the average surface temperature is about minus 235 degrees Celsius, it is currently summer in the southern hemisphere and winter in the northern. As Triton's southern hemisphere warms up, a thin layer of frozen nitrogen, methane, and [carbon monoxide](#) on Triton's surface sublimates into gas, thickening the icy atmosphere as the season progresses during Neptune's 165-year orbit around the Sun. A season on Triton lasts a little over 40 years, and Triton passed the southern summer solstice in 2000.

Based on the amount of gas measured, Lellouch and his colleagues estimate that Triton's atmospheric pressure may have risen by a factor of four compared to the measurements made by Voyager 2 in 1989, when it was still spring on the giant moon. The atmospheric pressure on Triton is now between 40 and 65 microbars — 20 000 times less than on Earth.

Carbon monoxide was known to be present as ice on the surface, but Lellouch and his team discovered that Triton's upper surface layer is enriched with carbon monoxide ice by about a factor of ten compared to the deeper layers, and that it is this upper "film" that feeds the atmosphere. While the majority of Triton's atmosphere is nitrogen (much like on Earth), the methane in the atmosphere, first detected by Voyager 2, and only now confirmed in this study from Earth, plays an important role as well. "Climate and atmospheric models of Triton have to be revisited now, now that we have found carbon monoxide and re-measured the methane," says co-author Catherine de Bergh.

Of Neptune's 13 moons, Triton is by far the largest, and, at 2700 kilometres in diameter (or three quarters the Earth's Moon), is the seventh largest moon in the whole Solar System. Since its discovery in

1846, Triton has fascinated astronomers thanks to its geologic activity, the many different types of surface ices, such as frozen nitrogen as well as water and dry ice (frozen carbon dioxide), and its unique retrograde motion.

Observing the atmosphere of Triton, which is roughly 30 times further from the Sun than Earth, is not easy. In the 1980s, astronomers theorised that the atmosphere on Neptune's moon might be as thick as that of Mars (7 millibars). It wasn't until Voyager 2 passed the planet in 1989 that the atmosphere of nitrogen and [methane](#), at an actual pressure of 14 microbars, 70 000 times less dense than the atmosphere on Earth, was measured. Since then, ground-based observations have been limited. Observations of stellar occultations (a phenomenon that occurs when a Solar System body passes in front of a star and blocks its light) indicated that Triton's surface pressure was increasing in the 1990's. It took the development of the Cryogenic High-Resolution Infrared Echelle Spectrograph (CRIRES) at the Very Large [Telescope](#) (VLT) to provide the team the chance to perform a far more detailed study of Triton's atmosphere. "We needed the sensitivity and capability of CRIRES to take very detailed spectra to look at the very tenuous atmosphere," says co-author Ulli Käufl. The observations are part of a campaign that also includes a study of Pluto [eso0908 - <http://www.eso.org/public/news/eso0908/>].

Pluto, often considered a cousin of Triton and with similar conditions, is receiving renewed interest in the light of the carbon monoxide discovery, and astronomers are racing to find this chemical on the even more distant dwarf planet.

This is just the first step for astronomers using CRIRES to understand the physics of distant bodies in the [Solar System](#). "We can now start monitoring the [atmosphere](#) and learn a lot about the seasonal evolution of Triton over decades," Lellouch says.

**More information:** This research was presented in a paper to appear in *Astronomy & Astrophysics* ("Detection of CO in Triton's atmosphere and the nature of surface-atmosphere interactions", by E. Lellouch et al.), reference [DOI:10.1051/0004-6361/201014339](https://doi.org/10.1051/0004-6361/201014339)

Provided by ESO

Citation: Triton's summer sky of methane and carbon monoxide (2010, April 7) retrieved 25 April 2024 from <https://phys.org/news/2010-04-triton-summer-sky-methane-carbon.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.