

Cassini Double Play: Enceladus and Titan

May 18 2010



On the left, Saturn's moon Enceladus is backlit by the sun, showing the fountain-like sources of the fine spray of material that towers over the south polar region. On the right, is a composite image of Titan. Image credit: NASA/JPL/SSI and NASA/JPL/University of Arizona

(PhysOrg.com) -- About a month and a half after its last double flyby, NASA's Cassini spacecraft will be turning another double play this week, visiting the geyser moon Enceladus and the hazy moon Titan. The alignment of the moons means that Cassini can catch glimpses of these two contrasting worlds within less than 48 hours, with no maneuver in between.

Cassini will make its closest approach to Enceladus late at night on May 17 Pacific time, which is in the early hours of May 18 UTC. The spacecraft will pass within about 435 kilometers (270 miles) of the moon's surface.

The main scientific goal at [Enceladus](#) will be to watch the sun play

peekaboo behind the water-rich plume emanating from the moon's south polar region. Scientists using the ultraviolet imaging spectrograph will be able to use the flickering light to measure whether there is molecular nitrogen in the plume. Ammonia has already been detected in the plume and scientists know heat can decompose ammonia into nitrogen molecules. Determining the amount of molecular nitrogen in the plume will give scientists clues about thermal processing in the moon's interior.

The second of Cassini's two flybys is an encounter with Titan. The closest approach will take place in the late evening May 19 Pacific time, which is in the early hours of May 20 UTC. The spacecraft will fly to within 1,400 kilometers (750 miles) of the surface.

Cassini will primarily be doing radio science during this pass to detect the subtle variations in the [gravitational tug](#) on the spacecraft by Titan, which is 25 percent larger in volume than the [planet Mercury](#). Analyzing the data will help scientists learn whether Titan has a liquid ocean under its surface and get a better picture of its internal structure. The [composite infrared spectrometer](#) will also get its southernmost pass for thermal data to fill out its temperature map of the smoggy moon.

Cassini has made four previous double flybys and one more is planned in the years ahead.

More information: More information on the Enceladus flyby, dubbed "E10," is available at: saturn.jpl.nasa.gov/mission/flybys/enceladus20100518/

More information on the Titan flyby, dubbed "T68," is available at: saturn.jpl.nasa.gov/mission/flybys/titan20100520/

Provided by JPL/NASA

Citation: Cassini Double Play: Enceladus and Titan (2010, May 18) retrieved 9 April 2024 from <https://phys.org/news/2010-05-cassini-enceladus-titan.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.