


# Revealing the metabolic activity of microbial communities

May 11 2010



A decimal place slope is a fast and precise method for quantifying  $^{13}\text{C}$  incorporation levels for detecting the metabolic activity of microbial species.  
 Credit: Stephan Boehme/UFZ

Microbial communities are performing important functions all around us - from the earth in our flowerpots to the human gut. Now researchers have developed a method for studying the metabolic functions of microbial communities in detail. It is now possible for the first time, thanks to a new algorithm developed at the UFZ (Germany), to use the incorporation of stable carbon isotopes into proteins to investigate natural remineralisation processes in much greater detail, to identify relevant key species and to study the way they interact in complex decomposition processes.

The new Protein-SIP technique makes it possible to measure carbon flux

in [microbial communities](#) very accurately, say researchers from the Helmholtz Centre for Environmental Research (UFZ), the Max Planck Institute for Infection Biology and the Universities of Oslo and Greifswald writing in *Molecular and Cellular Proteomics*.

Although in the past it was possible to identify species with metabolic activity using DNA or RNA analyses, the new method can also identify carbon flux and therefore food chains within a microbial community. This means that it is now possible to analyse the interaction between individual groups of micro-organisms within a community.

News headlines like "Uncovering the genetic secrets of intestinal bacteria" and "Hand bacteria used to catch criminals" show that microbiologists all over the world are currently working hard to explore the world of bacteria living on and in the human body. The scope of potential applications is huge and could range from forensic medicine and simpler medical diagnosis to entirely new treatments. However, simply identifying the genes is not enough, because bacteria do not live on their own, but in large communities. "It's like a city with lots of people. Imagine a fire breaks out. Normally, fire-fighters would deal with it, but if there are no fire-fighters around, other people have to step in to prevent disaster," explains Dr Ingo Fetzer of the UFZ. "But who is responsible for what within these microbial communities? This is an important question that scientists are only just starting to investigate." And it is not just [human gut](#) flora that are at issue. Microbes are tiny organisms that, unseen by the human eye, control all the major biological processes on earth - from the global carbon cycle to the remineralisation of organic material and the breakdown of harmful substances.

The number of species of higher organisms on the planet is estimated to be between five and 100 million. There are only vague conjectures about the number of species of micro-organisms. This means that researchers have to concentrate on just a few species. So how is it possible to

identify the key organisms within the microbial communities? In order to answer this question more easily, researchers at the Helmholtz Centre for Environmental Research combined the use of stable isotopes with protein measurements using mass spectrometry and bioinformatics. In the new method, microbial communities are fed a carbon source containing the heavy, non-radioactive isotope  $^{13}\text{C}$  as well as normal carbon,  $^{12}\text{C}$ . The two isotope masses differ by 1.0035 atomic mass units. Because they are stable isotopes, the method is also known as Stable Isotope Probing (SIP). Once the bacteria have consumed the isotope-marked substrate, the  $^{13}\text{C}$  atoms are incorporated into the bacterial proteins. The bacteria that make use of the substrate itself incorporate the  $^{13}\text{C}$  first. Other species of bacteria only make use of metabolites from the first group and incorporate less  $^{13}\text{C}$  into their proteins and do so later. For the analysis, the proteins of all the bacterial species from a sample are extracted and cut into specific fragments using the enzyme trypsin. The fragments are analysed using a mass spectrometer to determine the amino acid sequence of the peptides. When compared with a genome database, this reveals a peptide's origin, i.e. the bacterium it comes from. Peptides are protein fragments - organic compounds containing a number of amino acids. These consist primarily of carbon and nitrogen, which are two of the basic building blocks of all molecules within organisms and are therefore passed on even in mixed microbial cultures. In a second step, the researchers calculate the level of  $^{13}\text{C}$  incorporation. The  $^{13}\text{C}$  level then provides an elegant, direct and accurate measure of the metabolic activity of the species in question. "We first tested this key technology in 2008 in a joint project conducted by two UFZ departments to analyse the [metabolic activity](#) of one specific species of bacteria within a mixed culture. We have been studying the structure and function of the microbial communities involved in the breakdown harmful substances for years. But it is only with the advent of the new mass spectrometers and their more accurate measurements that we have been able to achieve a breakthrough in developing the method," says project coordinator Dr

Martin von Bergen from the Department of Proteomics.

Now it is possible to calculate the level of  $^{13}\text{C}$  incorporation into the peptides using the decimal places of the peptide masses. The researchers make use of the 0.0035 deviance in atomic mass units over and above the theoretically precise figure of 1.000 atomic mass units between  $^{12}\text{C}$  and  $^{13}\text{C}$ . Since there are more than 20 carbon atoms in a peptide, the decimal places are shifted over around 0.07 atomic mass units. Prof. Hauke Harms from the Department of Environmental Microbiology is very pleased with the new method: "Our new algorithm will make research work much easier in future. The method offers great potential for studying communities, which are at the heart of microbial ecology."

With support from the German Research Foundation (DFG) and the EU, researchers will now identify the key organisms in the breakdown of environmental pollutants such as benzene and polycyclic hydrocarbons in the absence of oxygen. "In conjunction with other techniques, Protein-SIP is a very good tool for investigating the food web involved in the breakdown of benzene, for example. Protein-SIP is already being used in projects with national and international partners to identify the metabolic activities of methane bacteria from oil deposits and the methane cycle in marine sediments," Dr Hans Richnow (Department of Isotope Biogeochemistry) adds. These projects are of relevance for securing energy supplies and conserving the quality of the environment.

The Protein-SIP method makes it possible to trace the carbon flux within mixed bacterial cultures. Other potential applications include the treatment of biofilms, such as those used in sewage works, and the optimisation of biogas generation processes and the analysis of the human intestine.. The next step for the Leipzig-based researchers is to examine the relationship between the [intestinal bacteria](#) of termites and earthworms and their host organisms.

The United Nations have declared 2010 as the 'International Year of Biodiversity'. The goal of this is to bring the issue of biodiversity with its many facets to the collective conscience of the public. With its expertise the UFZ contributes to investigating the consequences and causes of the loss of biodiversity as well as developing mitigation options.

**More information:** Publications:

Nico Jehmlich, Ingo Fetzer, Jana Seifert, Jens Mattow, Carsten Vogt, Hauke Harms, Bernd Thiede, Hans Hermann Richnow, Martin von Bergen, and Frank Schmidt (2010): Decimal place slope: a fast and precise method for quantifying <sup>13</sup>C incorporation levels for detecting the metabolic activity of microbial species Mol Cell Proteomics.

[dx.doi.org/10.1074/mcp.M900407-MCP200](http://dx.doi.org/10.1074/mcp.M900407-MCP200)

Jehmlich N, Schmidt F, Hartwich M, von Bergen M, Richnow HH, Vogt C. Incorporation of carbon and nitrogen atoms into proteins measured by protein-based stable isotope probing (Protein-SIP). Rapid Commun Mass Spectrom. 2008 Sep;22(18):2889-97.

[www3.interscience.wiley.com/jo ... /1/121387805/abstract](http://www3.interscience.wiley.com/jo.../1/121387805/abstract)

Jehmlich N, Schmidt F, Taubert M, Seifert J, von Bergen M, Richnow HH, Vogt C. Comparison of methods for simultaneous identification of bacterial species and determination of metabolic activity by protein-based stable isotope probing (Protein-SIP) experiments. Rapid Commun Mass Spectrom. 2009 Jun;23(12):1871-8.

[www3.interscience.wiley.com/jo ... /1/122385820/abstract](http://www3.interscience.wiley.com/jo.../1/122385820/abstract)

Jehmlich, N., Schmidt, F., von Bergen, M., Richnow, H.-H., Vogt, C. (2008): Protein-based stable isotope probing (Protein-SIP) reveals active species within anoxic mixed cultures. The ISME Journal 2, 1122-1133

[dx.doi.org/10.1038/ismej.2008.64](http://dx.doi.org/10.1038/ismej.2008.64)

Provided by Helmholtz Association of German Research Centres

Citation: Revealing the metabolic activity of microbial communities (2010, May 11) retrieved 19 April 2024 from <https://phys.org/news/2010-05-revealing-metabolic-microbial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.