

## **Report describes the physics of the 'bends'**

June 22 2010

As you go about your day-to-day activities, tiny bubbles of nitrogen come and go inside your tissues. This is not a problem unless you happen to experience large changes in ambient pressure, such as those encountered by scuba divers and astronauts. During large, fast pressure drops, these bubbles can grow and lead to decompression sickness, popularly known as "the bends."

A study in the <u>Journal of Chemical Physics</u>, which is published by the American Institute of Physics (AIP), may provide a physical basis for the existence of these bubbles, and could be useful in understanding <u>decompression sickness</u>.

A physiological model that accounts for these bubbles is needed both to protect against and to treat decompression sickness. There is a problem though. "These bubbles should not exist," says author Saul Goldman of the University of Guelph in Ontario, Canada.

Because they are believed to be composed mostly of nitrogen, while the surrounding atmosphere consists of both nitrogen and oxygen, the <u>pressure</u> of the bubbles should be less than that of the surrounding atmosphere. But if this were so, they would collapse.

"We need to account for their apparent continuous existence in tissues in spite of this putative pressure imbalance," says Goldman.

If, as is widely believed, decompression sickness is the result of the growth of pre-existing gas bubbles in tissues, those bubbles must be



sufficiently stable to have non-negligible half-lives. The proposed explanation involves modeling body tissues as soft elastic materials that have some degree of rigidity. Previous models have focused on bubble formation in simple liquids, which differ from elastic materials in having no rigidity.

Using the soft-elastic tissue model, Goldman finds pockets of reduced pressure in which nitrogen bubbles can form and have enough stability to account for a continuous presence of tiny bubbles that can expand when the ambient pressure drops. Tribonucleation, the phenomenon of formation of new gas bubbles when submerged surfaces separate rapidly, provides the physical mechanism for formation of new gas bubbles in solution. The rapid separation of adhering surfaces results in momentary negative pressures at the plane of separation. Therefore, while these tiny bubbles in elastic media are metastable, and do not last indefinitely, they are replaced periodically. According to this picture, tribonucleation is the source, and finite half-lives the sink, for the continuous generation and loss small <u>gas bubbles</u> in tissues.

**More information:** The article, "Free energy wells for small gas bubbles in soft deformable materials" by Saul Goldman was published on April 26,2010 in the Journal of Chemical Physics. See: jcp.aip.org/jcpsa6/v132/i16/p164509\_s1

## Provided by American Institute of Physics

Citation: Report describes the physics of the 'bends' (2010, June 22) retrieved 3 May 2024 from <u>https://phys.org/news/2010-06-physics.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.