

## Scientists get a look at the birth of the Milky Way

June 21 2010



An image of the globular star cluster M80 (NGC 6093) made using the Hubble Space Telescope. M80 contains hundreds of thousands of stars and is one of 147 globular clusters known to be associated with the Milky Way. Credit: The Hubble Heritage Team / AURA / STScI / NASA

(PhysOrg.com) -- For the first time, a team of astronomers has succeeded in investigating the earliest phases of the evolutionary history of our home Galaxy, the Milky Way. The scientists, from the Argelander Institute for Astronomy at Bonn University and the Max-Planck Institute for Radioastronomy in Bonn, deduce that the early Galaxy went from smooth to clumpy in just a few hundred million years. The team publish



their results in the journal *Monthly Notices of the Royal Astronomical Society*.

Led by Professor Dr. Pavel Kroupa, the researchers looked at the spherical groups of stars (globular clusters) that lie in the halo of the Milky Way, outside the more familiar <u>spiral arms</u> where the Sun is found. They each contain hundreds of thousands of stars and are thought to have formed at the same time as the 'proto-Galaxy' that eventually evolved into the Galaxy we see today.

Globular star clusters can be thought of as fossils from the earliest period of the history of the Galaxy and the <u>astronomers</u> found that they left a hint of the conditions under which they formed. The stars of the clusters condensed out of a cloud of molecular gas (relatively cool hydrogen), not all of which was used up in their formation. The residual gas was expelled by the radiation and winds coming from the freshly hatched population of stars.

"Due to this ejection of gas, the globular clusters expanded and thereby lost the stars that formed at their boundaries. This means that the present shape of the clusters was directly influenced by what happened in the early days of their existence", explains Michael Marks, PhD student of Professor Kroupa and lead author on the new paper.

The clusters were also shaped by the forming Milky Way and the Bonn scientists calculated exactly how the proto-Galaxy affected its smaller neighbours. Their results show that the gravitational forces exerted on the star clusters by the proto-Milky Way appear to increase with the metal content of their member stars (in astronomy 'metals' in stars are elements heavier than helium).

"The amount of e.g. iron in a star is therefore an age indicator. The more recently a star cluster was born, the higher the proportion of heavy



elements it contains", adds Marks. But since the globular clusters are more or less the same age, these age differences can't be large. In order to explain the variation in the forces exerted on different globular clusters, the structure of the Milky Way had to change rapidly within a short time.

The giant gas cloud from which the <u>Milky Way</u> formed had to evolve from an overall smooth structure into a clumpy object in less than a few hundred million years in order to increase the strength of the forces significantly. This timespan corresponds to the astronomically short duration in which the proto-galaxy-sized gas cloud collapsed under its own gravity. In parallel, the globular clusters formed successively within the collapsing cloud. The material from which the somewhat younger globular clusters formed and which according to the results of this investigation felt stronger attractive forces, was previously enriched with heavy elements by fast-evolving stars in the older clusters.

Prof. Kroupa summarises their results. "In this picture we can elegantly combine the observational and theoretical results and understand why later forming, more metal-rich clusters experienced stronger force fields. On the back of this work, for the first time we have a detailed insight into the earliest evolutionary history of our Galaxy".

**More information:** The new research appears in "Initial conditions for globular clusters and assembly of the old globular cluster population of the Milky Way", Marks M., Kroupa P., Monthly Notices of the Royal Astronomical Society, in press. A preprint of this paper can be seen at <u>arxiv.org/abs/1004.2255</u>

Provided by Royal Astronomical Society



Citation: Scientists get a look at the birth of the Milky Way (2010, June 21) retrieved 2 May 2024 from <u>https://phys.org/news/2010-06-scientists-birth-milky.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.