

## **Tracking coral larvae to understand Hawai'i reef health**

June 9 2010



This underwater photograph shows larvae being released into the water column from reef-building coral spawning off Maui the night of June 30, 2003. The white larvae are about 0.5 to 1.5 mm across. This is the same coral species being studied in Maunalua Bay, Oahu, this summer. Credit: Photograph copyright 2003 by John Gorman, Maui Ocean Center. Permission is granted to use this photo to accompany stories resulting from this news release.

Scientists can predict almost to the hour when the reef-building "rice coral" off Oahu will spawn, but no one knows where the resulting floating coral larvae go.

From June 11-16, 2010, the U.S. Geological Survey, the University of Hawai'i at Mānoa's Kewalo Marine Laboratory, and Malama Maunalua will use satellite-tracked drifters to track the coral larvae's dispersal along O'ahu's south shore in an effort to better understand why certain



reefs in Maunalua Bay are doing well and others are doing poorly.

The next coral spawning event will begin after the next new moon, which is June 12. The scientists will use underwater instruments to monitor tides, waves, currents, temperature, salinity, chlorophyll, and water clarity throughout the bay. They will release satellite-tracked drifters to float along with the buoyant coral eggs and larvae and monitor the drifters' positions all night to determine if they are being transported into, retained in, or dispersed out of Maunalua Bay.

During the experiment, boaters and swimmers should keep an eye out for, and avoid the small (8-inch diameter) orange drifters and some small yellow floats marking instrument packages, all with "USGS" markings.

Maunalua Bay has been degraded by polluted runoff and sediment, invasive <u>algae</u>, and unsustainable harvesting. Reduced water and bottom quality tied to watershed discharges and invasive algae affect not only the corals in the bay, but the ability of <u>larvae</u> from elsewhere to replenish depleted populations. Community-based efforts to remove invasive algae and reduce runoff and sedimentation will ultimately lead to improved coral and fish resources. Recovery of the Bay also depends in part on the reproductive health of the local and adjacent coral reefs as sources of <u>coral</u> seed.

Because they are living organisms, corals are born and die. If new corals are not produced and able to settle, replacing those that die from both natural and human-induced causes, the reefs will eventually disappear, along with the other associated marine resources. Once scientists understand the circulation, larval dispersal patterns, and "connectivity" between reefs, managers can identify where recovery efforts should be focused.



## Provided by University of Hawaii at Manoa

Citation: Tracking coral larvae to understand Hawai'i reef health (2010, June 9) retrieved 2 May 2024 from <u>https://phys.org/news/2010-06-tracking-coral-larvae-hawaii-reef.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.