

Fungi's genetic sabotage in wheat discovered

July 13 2010

Using molecular techniques, Agricultural Research Service (ARS) and collaborating scientists have shown how the subversion of a single gene in wheat by two fungal foes triggers a kind of cellular suicide in the grain crop's leaves.

Fortunately, the team has also developed DNA molecular markers that can be used to rapidly screen commercial cultivars for the gene, *Tsn1*, so it can be eliminated by [selective breeding](#). This, in turn, would deprive the fungi of their primary means of killing off leaf tissue to feed and grow, explains Justin Faris, a [plant geneticist](#) with the ARS Cereal Crops Research Unit in Fargo, N.D.

The fungi—*Pyrenophora tritici-repentis* (also known as tan spot) and *Stagonospora nodorum* (leaf blotch)—are often partners in crime, occurring in the same crop fields and producing the same toxin, ToxA, to induce a *Tsn1*-controlled response in wheat called programmed cell death (PCD). Normally, PCD protects plants by confining invading pathogens in dead cells. However, the strategy doesn't work against the ToxA fungi because they're "necrotrophs," pathogens that feed on dead tissue.

To better understand this genetic trickery, Faris led a team of scientists from seven different research organizations in isolating, sequencing and cloning the DNA sequence for *Tsn1* from cultivated wheat and its wild relatives. Based on their analysis, the researchers concluded that modern-day wheat inherited *Tsn1* from goatgrass. They figure this happened after a goatgrass gene for the [enzyme protein](#) kinase fused with another

gene, NB-LRR, which probably conferred resistance to biotrophs, pathogens that feed on living tissue.

Interestingly, Tsn1 is controlled by wheat's [circadian clock](#), and only initiates PCD in response to ToxA during daylight hours. At night, Tsn1 shuts down and "ignores" ToxA, suggesting the toxin may indirectly interfere with the plant's photosynthesis.

More information: The team, which includes researchers from North Dakota State University-Fargo and the Australian Centre for Necrotrophic Fungal Pathogens-Murdoch among others, reported its findings online this week in the *Proceedings of the National Academy of Sciences*.

Provided by United States Department of Agriculture

Citation: Fungi's genetic sabotage in wheat discovered (2010, July 13) retrieved 23 April 2024 from <https://phys.org/news/2010-07-fungi-genetic-sabotage-wheat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.