

La Nina developing, could mean more hurricanes

July 8 2010, By RANDOLPH E. SCHMID , AP Science Writer

(AP) -- The climate phenomenon known as La Nina appears to be developing, threatening more bad news in the efforts to clean up the oil spill in the Gulf of Mexico.

When a La Nina occurs there tend to be more hurricanes than normal in the Atlantic and Caribbean regions, which include the <u>Gulf of Mexico</u>.

The federal Climate Prediction Center said Thursday that La Nina conditions are likely to develop in July and August.

La Nina is marked by an unusual cooling of the sea surface in the tropical Pacific Ocean. Water temperatures in that area can affect air pressure and winds, resulting in changes in the weather in many parts of the world.

In a La Nina, <u>wind shear</u> is increased over the Pacific and reduced over the Atlantic. Wind shear is the difference in strength of winds at low levels compared to higher level winds.

A strong wind shear reduces hurricanes by breaking up their ability to rise into the air, while less shear means they can climb and strengthen.

Thus, the Climate Prediction Center notes, "there tend to be more Atlantic hurricanes during La Nina because of this expanded area of low vertical wind shear."



In addition, during a La Nina "more hurricanes form in the deep tropics from African easterly waves. These systems have a much greater likelihood of becoming major hurricanes, and of eventually threatening the U.S. and Caribbean islands," according to the center, a part of the National Oceanic and Atmospheric Administration.

The climate center's current <u>hurricane</u> forecast for this season is for 14 to 23 named storms of which 8 to 14 are expected to be hurricanes and 3 to 7 major hurricanes.

The center noted that during June, sea surface temperature continued to decrease across the equatorial Pacific Ocean, with cool areas expanding across the central and eastern Pacific. In addition, increased rainfall persisted over Indonesia, while the area of reduced rain expanded westward over the western and central equatorial Pacific.

Combined with changes in the winds over the Pacific "these oceanic and atmospheric anomalies reflect developing La Nina conditions" which are likely to continue through early 2011, CPC said.

The last <u>La Nina</u> occurred from the fall of 2007 to the spring of 2008. The opposite mode, El Nino, with warm Pacific conditions, has been in place since the spring of 2009.

More information: <u>http://www.cpc.ncep.noaa.gov/</u>

©2010 The Associated Press. All rights reserved. This material may not be published, broadcast, rewritten or redistributed.

Citation: La Nina developing, could mean more hurricanes (2010, July 8) retrieved 15 May 2024 from <u>https://phys.org/news/2010-07-la-nina-hurricanes.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.