

# 'Lab on a Chip' Detects Human, Agricultural Contaminants

July 19 2010, By Susan McGinley

---



The first prototype of lab-on-a-chip system for detecting microbes in water, food and air. (Photo credit: UA Biosensors Lab)

(PhysOrg.com) -- The UA's Jeong-Yeol Yoon is using glass-slide laboratories to detect E. coli in water and vegetables and to monitor disease in livestock.

Detecting water and foodborne contaminants usually involves collecting a sample, sending it to a laboratory and waiting for it to be filtered, incubated, tested and identified under a microscope. If a critical infection is suspected, say for E. coli, the pathogen may already have multiplied and spread before the report arrives days later.

A series of "[lab on a chip](#)" applications in development at the University

of Arizona can identify [pathogens](#) in minutes rather than days, using a simple device (which may be attached to a faucet) that can deliver results locally.

The degree of accuracy is three orders of magnitude greater than for conventional real-time or rapid tests, and the method can be used to test lower concentrations of pathogens.

Invented by chemical and electrical engineers in the late 1990s, the lab on a chip, or LOC, concept is based on the integrated circuit, where encapsulated wires and circuits are integrated into a [semiconductor chip](#) with electricity flowing through it.

"In lab on a chip, instead of using electricity, the liquid flows," said Jeong-Yeol Yoon, an assistant professor in the UA department of agricultural and biosystems engineering and a member of the BIO5 Institute.

When Yoon was hired at the UA in 2004, he began to test the technology on water after noticing that attempts by other scientists to test blood with the LOC failed.

"I realized lab on a chip needed to start with the simplest case. Water is easy and very dilute compared to blood," Yoon said. "When you run the test, there are almost no sizeable substances except the pathogens. It may detect a single cell."

The LOC is a small glass laboratory slide filled with nanoparticles that adhere to pathogens applied in a sample drop of water.

Yoon uses micro and nanoparticles from 1,000 nanometers down to 10 nanometers to scatter light and strengthen sensitivity. He combines them with antibodies to the target pathogens, leading to particle agglutination

and changing the scattering signal.

"An antibody with nanoparticles gives a stronger signal," Yoon said. "Others use fluorescent dye; we use nanoparticles as a substitute for dye."

Working with graduate and undergraduate students in his Biosensors Laboratory on campus, Yoon has designed a test that can detect pathogens - E. coli and potentially Cryptosporidium - in drinking water networks, irrigation systems, or wastewater recycling facilities.

The prototype is finished but he is now devising different ways to attach the test to water systems, manipulate the [nanoparticles](#) and read and deliver the results by computer.

Yoon also has funding from a government agency in his native South Korea to develop tests for livestock diseases in that country.

These include LOC biosensors for bovine viral diarrhea, a bacterial disease that can reduce the quality of the milk and meat, and porcine reproductive and respiratory syndrome, a common viral disease that affects both the amount and the quality of the pork produced.

Yoon is also monitoring the viral pathogens for avian flu and H1N1 flu in South Korean livestock environments.

"Water supplies are affected first because the animals touch their tongues to the faucets," he said. "We're installing an LOC system on that water supply and also testing their wastewater."

The goal is to meet increased consumer demand for choice and premium grades of meat, and for higher quality milk, by detecting and treating diseases earlier.

Yoon is also testing air samples from livestock housing for detecting such viral pathogens. The air sample tests show about the same sensitivity and specificity as those with water samples.

To make vegetables safer, Yoon is developing a test for bacterial contamination - specifically E. coli and Salmonella - in fresh lettuce and spinach.

"We collect lettuce, grind it up, turn it into vegetable juice and then load it onto a lab on a chip and read the results," he said. "It's the most complicated application so far, but it's working well. We're detecting pathogens not from water but directly from the vegetable tissues."

The effort is funded by Desert Tech Investors.

Yoon's ultimate step is a human biomedical application: creating an LOC test for the H1N1 virus. For this, he is using newer "droplet" technology, where droplets on nano-structured surfaces can be manipulated using a control pad - the current prototype in the Biosensors Lab is attached to a Nintendo gaming pad.

"You can choose the reagent droplets you want (either DNA or antibody), move the sample droplet to them, and mix, rotate and/or heat them depending on the nature of the sample," Yoon said. "It's more expensive because you can change the protocol. The circuit memorizes the previous movements for 100 different samples."

The rapid-response test can be done in real time and will be commercially available in about five years.

Provided by University of Arizona

Citation: 'Lab on a Chip' Detects Human, Agricultural Contaminants (2010, July 19) retrieved 19 July 2024 from <https://phys.org/news/2010-07-lab-chip-human-agricultural-contaminants.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.