

By 'putting a ring on it,' microparticles can be captured

July 20 2010



This is a schematic illustration of a particle revolving around a silicon micro-ring resonator, propelled by optical forces. Credit: Courtesy of Ken Crozier, Harvard School of Engineering and Applied Sciences.

To trap and hold tiny microparticles, engineers at Harvard have "put a ring on it," using a silicon-based circular resonator to confine particles stably for up to several minutes.

The advance, published in the June 14 issue of <u>Nano Letters</u>, could one day lead to the ability to direct, deliver, and store <u>nanoparticles</u> and <u>biomolecules</u> on all-optical chips.

"We demonstrated the power of what we call resonant cavity trapping, where a particle is guided along a small waveguide and then pulled onto a micro-ring resonator," explains Kenneth Crozier, an Associate Professor of Electrical Engineering at the Harvard School of Engineering and Applied Sciences (SEAS) who directed the research. "Once on the ring, optical forces prevent it from escaping, and cause it



to revolve around it."

The process looks similar to what you see in liquid motion toys, where tiny beads of colored drops run along plastic tracks—but on much smaller scale and with different physical mechanisms. The rings have radii of a mere 5 to 10 micrometers and are built using <u>electron beam</u> <u>lithography</u> and reactive ion etching.

Specifically, <u>laser light</u> is focused into a waveguide. Optical forces cause a particle to be drawn down toward the waveguide, and pushed along it. When the particle approaches a ring fabricated close to the waveguide, it is pulled from the waveguide to the ring by optical forces. The particle then circulates around the ring, propelled by optical forces at velocities of several hundred micrometers-per-second.

While using planar ring resonators to trap particles is not new, Crozier and his colleagues offered a new and more thorough analysis of the technique. In particular, they showed that using the silicon ring results in optical force enhancement (5 to 8 times versus the straight waveguide).



This is a scanning electron micrograph (SEM) of the silicon micro-ring resonator



(radius: 5 micron) coupled to waveguide. Credit: Courtesy of Kenneth Crozier, Harvard School of Engineering and Applied Sciences

"Excitingly, particle-tracking measurements with a high speed camera reveal that the large transverse forces stably localize the particle so that the standard deviation in its trajectory, compared to a circle, is as small as 50 nm," says Crozier. "This represents a very tight localization over a comparatively large distance."

The ultimate aim is to develop and demonstrate fully all-optical on chip manipulation that offers a way to guide, store, and deliver both biological and artificial particles.

Provided by Harvard University

Citation: By 'putting a ring on it,' microparticles can be captured (2010, July 20) retrieved 2 May 2024 from <u>https://phys.org/news/2010-07-microparticles-captured.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.