

Research shows it is not about more sunlight and higher temperatures

July 20 2010



A recent scientific study of insect diversity suggests that it's seasonality, not heat or light, that drives biodiversity. Photo by S. Bruce Archibald

The tropics owe their stunning biodiversity to consistent year-round temperatures, not higher temperatures or more sunlight, according to a novel survey of insect diversity at different latitudes and at different points in the planet's history.

The finding, presented this week (July 20) in the journal *Paleobiology* by researchers from Harvard University, Simon Fraser University, and Brandon University, may finally answer a question that has dogged scientists for centuries.

It also suggests, intriguingly, that the world is likely far less diverse today than it was tens of millions of years ago, when the entire Earth had

consistent year-round temperatures, much like the modern tropics.

“The latitudinal diversity gradient has been recognized for 150 years as one of the most general observations in nature, and has produced more explanatory hypotheses than nearly any other observation,” says co-author Brian D. Farrell, professor of biology at Harvard. “We show that when most of today’s organisms were diversifying, up through the Eocene [epoch], the world lacked pronounced [seasonality](#), more like today’s tropics, even in areas where the temperature was low.”

“It appears it’s not the heat of the tropics that promotes diversity; it’s the newer seasons of the temperate zone that depress diversity.”

Scientists’ explanations for tropical biodiversity have tended to focus on the greater heat and light found closer to the equator, and to a lesser extent on the low seasonality of the tropics, where average temperature in the hottest and coolest months may vary by only a few degrees.

“These factors tend to change together as you travel away from the equator to-ward the poles, leaving it difficult to separate their individual effects on diversity,” says lead author S. Bruce Archibald, a research associate at Simon Fraser University, Harvard’s Museum of Comparative Zoology, and the Royal British Columbia Museum. Archibald conducted the research for his doctoral dissertation at Harvard, where Farrell was his adviser.

Farrell, Archibald, and colleagues engaged in a kind of time travel, invoking the fossil record to solve the conundrum of biodiversity. They compared modern insect diversity at the Harvard Forest in Petersham, Mass., and in a Costa Rican jungle against that seen at the 52.9 million-year-old McAbee fossil bed in British Columbia, noted for its exceptionally well preserved insects. At the time the McAbee fossils were created, Earth’s climate was far less seasonal at all latitudes,

allowing tropical species such as palm trees and crocodiles to live in what is now the high Arctic.

The scientists' discovery that the ancient Canadian site's insect diversity mirrors that of the modern Costa Rican jungle, despite a marked difference in latitude, suggests that it's seasonality, not heat or light, that drives biodiversity.

“Planet Earth and life have co-evolved for well over a billion years, and this is yet additional evidence of the consequent intertwining of global physical and biological phenomena,” Farrell says.

Provided by Harvard University

Citation: Research shows it is not about more sunlight and higher temperatures (2010, July 20) retrieved 23 April 2024 from <https://phys.org/news/2010-07-sunlight-higher-temperatures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.