

Spinal gap of barbeled dragonfishes mystery solved

August 17 2010



Stained specimen of the dragonfish, *Chauliodus*, shows the gap in its backbone, which is due to missing vertebrae. There is also another explanation for the gap in other dragonfishes. © Nalani Schnell

(PhysOrg.com) -- The mystery of why a group of deep sea fishes has a gap between their skull and spine has been solved by an international team of scientists, including those at the Natural History Museum.

Large sharp teeth, a thread-like skin extension on their chins called a barbel, and protruding jaws make barbeled dragonfishes striking enough.

But they also appear to have a number of vertebrae missing that would normally connect their skulls to their vertebral column, or backbone, an observation that has puzzled scientists for many years.



Deep sea dragonfish Eustomias with its striking barbel. This is one of the dragonfishes that was found to have vertebrae missing. © Nalani Schnell

There are 28 genera (major groups) of these fishes, representing more than 270 species, and all have this characteristic gap. But are missing vertebrae always the cause?

Most detailed study

Dr Nalani Schnell from the University of Tübingen, Germany, along with [Natural History Museum](#)'s ichthyologist (fish expert) Dr Ralf Britz and Dr David Johnson from the Smithsonian Institution in Washington DC, USA, decided to investigate.


The team carried out the most detailed and comprehensive study so far of this group.

They stained the fish specimens so that nerves and bones were shown in different colours, producing images (such as the one at the top) and revealing clues about the mystery gap.

The team studied some specimens from the Museum's fish collection, which includes more than 2000 dragonfish specimens.

Vertebrae lost during evolution

The team found vertebrae missing in members of only 2 of the genera of barbeled dragonfishes, *Chauliodus*, and *Eustomias*, and in the species *Leptostomias gladiator*. This loss occurred as they evolved.



Stained dragonfishes showing gaps in their vertebral column, top to bottom, *Flagellostomias boureei*, *Grammatostomias dentatus* and *Malacosteus niger*. © Nalani Schnell

Usually, fish backbones develop from front to back, (top to bottom in humans). However, in barbeled dragonfishes this is not the case, making

the evolutionary loss of front vertebrae more likely.

Dr Britz explains, ‘In stomiids [barbeled dragonfishes are in the Family Stomiidae], vertebrae develop in an unusual fashion from back to front, which facilitated the loss of front vertebrae. They have failed to form in development leaving a gap between the skull and backbone.

‘This results in them having a more flexible head and makes them unique among all bony fishes in this respect.’

Gap due to long notochord

The gap in the backbone of the other barbeled dragonfishes studied are not caused by missing [vertebrae](#) but by a lengthening of the notochord. The notochord is the primitive support structure for the backbone, a flexible rod-like structure that in higher vertebrates is incorporated into or replaced by the vertebral column.

Barbeled dragonfishes

Barbeled dragonfishes are deep sea ray-finned fish. Most lack scales and they have a series of light-emitting organs called photophores.

The barbel that hangs from the lower jaw carries one of these photophores at its tip. It emits a blue light and is thought to help attract prey.

This research is published online and in the July/August printed issue of *Journal of Morphology*.

Provided by American Museum of Natural History

Citation: Spinal gap of barbeled dragonfishes mystery solved (2010, August 17) retrieved 11 May 2024 from <https://phys.org/news/2010-08-spinal-gap-barbeled-dragonfishes-mystery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.