


# The art of dividing: Researchers decode function and protein content of the centrosome

September 6 2010

---



Left: During cell division, chromosomes (red) are distributed evenly by thread-like structures (microtubules) emanating from the centrosome (green). Right: Inactivation of a centrosomal protein causes abnormal organisation of the mitotic spindle microtubules and a faulty distribution of chromosomes. Image: Bodo Lange

A basic requirement for growth and life of a multicellular organism is the ability of its cells to divide. A protein complex, the so-called centrosome, plays a major role during cell division. Scientists at the Max Planck Institute for Molecular Genetics in Berlin, together with colleagues at the German Cancer Research Center in Heidelberg and at the Leibniz Institute for Age Research - Fritz Lipmann Institute in Jena have investigated the functions of the different centrosomal components. The researchers now present the centrosome's components and their functions. Their work extends our knowledge of regulation of cell

division and opens the door to new investigations into cancer development. (*EMBO Journal*, September 3rd 2010)

During cell division chromosomes duplicate and are then distributed among the daughter cells. This distribution is organized by the [centrosome](#), a protein complex made up of several hundred different proteins. In cancer cells, the centrosome often assumes an unnatural shape or is present in uncontrolled numbers. The reasons for this were previously largely unknown.

As part of their research, the scientists examined centrosomes of the fruit fly *Drosophila* as well as those from human cells. "The fruit fly is a terrific system for investigating the centrosome, because the basic mechanisms of cell division are very similar between fly and human", explains Bodo Lange, the research group leader at the Max Planck Institute for Molecular Genetics. The group isolated centrosomes from the eggs of [fruit flies](#) and, using mass spectrometry methods, identified more than 250 different proteins making up this complex. These components were then subjected to targeted inactivation through [RNA interference](#) (RNAi), to discover their role in the structure of the centrosome and in chromosome distribution. The scientists were able to determine the protein functions quantitatively through use of state-of-the-art automatic and robotic microscopes.

They found a whole series of proteins responsible for the separation of chromosomes, number of centrosomes and their structure. As these characteristics are often disrupted in cancer cells, the researchers believe their findings will have a significant impact on the understanding of cell division and the development of cancerous diseases.

The work of these scientists has brought new insight into the abnormalities seen in cancer cells. "Based on our findings, we hope to be able to unravel regulatory networks in the future, which will help to

target and interfere with the division of [cancer cells](#)."

**More information:** Proteomic and functional analysis of the *Drosophila* centrosome. *EMBO Journal*, online publication 03.09.2010, [doi:10.1038/emboj.2010.210](https://doi.org/10.1038/emboj.2010.210)

Provided by Max-Planck-Gesellschaft

Citation: The art of dividing: Researchers decode function and protein content of the centrosome (2010, September 6) retrieved 10 April 2024 from <https://phys.org/news/2010-09-art-decode-function-protein-content.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.