


Early sunflower family fossil found in South America

September 28 2010, by Lin Edwards



Fossil Asteraceae. Image credit: *Science*, DOI:10.1126/science.1193108

(PhysOrg.com) -- A beautifully preserved fossil identified as being of an early relative of the Asteraceae, or aster, family nearly 50 million years old suggests the plant family, which has now colonized much of the planet, originated in South America after Gondwana separated, forming South America, Australia, Africa, Antarctica and India.

Researcher Dr. Viviana Barreda and colleagues of the Argentinian Museum of National Sciences in Buenos Aires found the perfectly preserved [fossil](#) of two flowering heads in the north-west Patagonia region, in rocks along the river Picheleufu in what is now north-western Argentina, and dated it to 47.5 million years old. At the time the region

was sub-tropical, with average temperatures of around 19°C.

Fossils of the Asteraceae are rare and most often only fossilized pollen grains are found, but the Patagonian fossil features two complete flowers complete with winged seeds, petals and the stems.

The most striking feature of the fossil, which resembles a painting of sunflowers by 19th century Dutch painter Van Gogh, is a dense [flower head](#) (capitulum) like that forming the central part of a [sunflower](#) and providing a strong attraction for [pollinators](#).

According to Barreda, the finding and dating of the fossil suggest the ancestor of the Asteraceae probably originated in southern Gondwana, in a region that is now part of southern South America, before effective dispersal barriers were established within this massive [supercontinent](#). It had been thought that the aster family originated from a common ancestor of the Asteraceae, Goodeniaceae and Calyceraceae families in once-tropical Antarctica, migrating to Australia and South America as Antarctica cooled. Then, between 56 and 23 million years ago this [common ancestor](#) in South America branched into the Asteraceae and Calyceraceae. The new finding provides confirmation for this hypothesis.

Dr. Tod Stuessy, a botanist with the University of Vienna, who wrote an accompanying article to Barreda's paper, said the fossil was clear evidence of the existence of the sunflower family at its early stages of diversification.

Stuessy said little is known about the origins of the sunflower family and there is much still to learn about its evolution and how it colonized the rest of the planet so quickly or how the members of the family became so “incredibly diverse.”

The Asteraceae family is the most diverse family of flowering plants, and includes sunflowers, daisies, chrysanthemums, lettuce, and dandelions among its members. The family includes 20-30,000 species, and representatives are found on every continent with the exception of Antarctica.

The paper and article describing the results of two years of work on the fossil are published in the journal *Science*.

More information: Eocene Patagonia Fossils of the Daisy Family, V. D. Barreda, *Science* 24 September 2010: Vol. 329. no. 5999, p. 1621.
[DOI:10.1126/science.1193108](https://doi.org/10.1126/science.1193108)

© 2010 PhysOrg.com

Citation: Early sunflower family fossil found in South America (2010, September 28) retrieved 19 April 2024 from <https://phys.org/news/2010-09-early-sunflower-family-fossil-south.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.