


# Mars methane lasts less than a year

September 21 2010

---



Map of methane concentrations in Autumn (first martian year observed) overlaid on true colour map of Mars. Credit: NASA/Universita del Salento


A new study indicates that methane in the atmosphere of Mars lasts less than a year. Methane is replenished from localized sources that show seasonal and annual variations. This pattern of methane production raises questions as to whether the methane comes from geological activity - or biological processes.

Methane in the atmosphere of [Mars](#) lasts less than a year, according to a study by Italian scientists. Sergio Fonti (Universita del Salento) and Giuseppe Marzo (NASA Ames) have used observations from NASA's Mars Global Surveyor spacecraft to track the evolution of the gas over three martian years. They presented their results at the European [Planetary Science](#) Congress in Rome on Tuesday 21st September.

“Only small amounts of [methane](#) are present in the [martian atmosphere](#), coming from very localized sources. We've looked at changes in concentrations of the gas and found that there are seasonal and also

annual variations. The source of the methane could be geological activity or it could be biological -- we can't tell at this point. However, it appears that the upper limit for methane lifetime is less than a year in the martian atmosphere," said Fonti.

Levels of methane are highest in autumn in the northern hemisphere, with localized peaks of 70 parts per billion, although methane can be detected across most of the planet at this time of year. There is a sharp decrease in winter, with only a faint band between 40 and 50 degrees north. Concentrations start to build again in spring and rise more rapidly in summer, spreading across the planet.



Top: Map of methane concentrations in Autumn (first martian year observed). Peak emissions fall over Tharsis (home to the Solar System's largest volcano, Olympus Mons), the Arabia Terra plains and the Elysium region, also the site of volcanos. Bottom: True color map of Mars. Credit: NASA/Universita del Salento

“One of the interesting things that we’ve found is that in summer, although the general distribution pattern is much the same as in autumn, there are actually higher levels of methane in the southern hemisphere.


This could be because of the natural circulation occurring in the atmosphere, but has to be confirmed by appropriate [computer simulations](#),” said Fonti.

There are three regions in the northern hemisphere where methane concentrations are systematically higher: Tharsis and Elysium, the two main volcano provinces, and Arabia Terrae, which has high levels of underground water ice. Levels are highest over Tharsis, where geological processes, including magmatism, hydrothermal and geothermal activity could be ongoing.

“It’s evident that the highest concentrations are associated with the warmest seasons and locations where there are favorable geological -- and hence biological -- conditions such as geothermal activity and strong hydration. The higher energy available in summer could trigger the release of gases from geological processes or outbreaks of biological activity,” said Fonti.

The mechanisms for removing methane from the atmosphere are also not clear. Photochemical processes would not break down the gas quickly enough to match observations. However, wind driven processes can add strong oxidizers to the atmosphere, such as the highly reactive salt perchlorate, which could soak up methane much more rapidly.

Martian years are nearly twice as long as Earth years. The team used observations from the Thermal Emission Spectrometer (TES) on Mars Global Surveyor between July 1999 and October 2004, which corresponds to three martian years the team studied one of the characteristic spectral features of methane in nearly 3 million TES observations, averaging data together to eliminate noise.



This image shows concentrations of Methane discovered on Mars. Credit: NASA

“Our study is the first time that data from an orbiting spectrometer has been used to monitor methane over an extended period. The huge TES dataset has allowed us to follow the methane cycle in the martian atmosphere with unprecedented accuracy and completeness. Our observations will be very useful in constraining the origins and significance of martian methane,” said Fonti.

Methane was first detected in the martian atmosphere by ground based telescopes in 2003 and confirmed a year later by ESA’s Mars Express spacecraft. Last year, observations using ground based telescopes showed the first evidence of a seasonal cycle.

The [atmosphere](#) on Mars consists of 95% carbon dioxide, 3% nitrogen, 1.6% argon, and contains traces of oxygen and water, as well as methane.

Source: European Planetary Science Congress

Citation: Mars methane lasts less than a year (2010, September 21) retrieved 28 April 2024 from <https://phys.org/news/2010-09-mars-methane-year.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.