

## NIST ultraviolet source helps NASA spacecraft measure the origins of space weather

September 1 2010



NIST's unique "sliding spark source" (inside the glass tubing) feeds ultraviolet light into NASA's Solar Ultraviolet Magnetograph Investigation instrument, designed to measure magnetic fields on the sun. Credit: Reader/NIST

With a brilliant, finely tuned spark of ultraviolet (UV) light, a physicist at the National Institute of Standards and Technology helped NASA scientists successfully position a crucial UV sensor inside a space-borne instrument to observe a "hidden" layer of the Sun where violent space weather can originate.

Dark spots on the Sun release particles and electromagnetic fields into space. As these particles and fields pass through the Sun's "transition region," 5,000 kilometers above the surface, they can gather



considerable steam, resulting in violent episodes of "<u>space weather</u>" that can damage Earth-orbiting satellites and disrupt electronic communications.

The powerful magnetic fields in the transition region can be studied indirectly, by observing the UV light emanating from that region. The fields slightly shift the colors (wavelengths) of UV light released by charged atoms (ions) in their vicinity. Measuring how much these wavelengths shift can yield information on the magnetic field's strength.

The catch is you can't do it from Earth, where the atmosphere absorbs the UV light, so a team at NASA Marshall Space Flight Center in Huntsville, Ala., constructed a rocket-borne instrument, known as the Solar Ultraviolet Magnetograph Investigation (SUMI), designed to take pictures of these magnetic fields from space.

SUMI observes shifts in the well-known wavelengths of UV light emitted by magnesium and carbon ions caught in the magnetic fields of the transition region. The instrument's optics break down the incoming UV light into a spectrum of individual wavelengths and fans them out, much as a prism fans out white light into a rainbow. The trick is in knowing precisely which <u>wavelength</u> falls where in the instrument and adjusting it so that the desired wavelengths land on the instrument's detectors.

"The problem is that SUMI's detectors are small, so they don't capture a wide range of wavelengths," says NIST physicist Joseph Reader. "The issue becomes how to align the complicated optics in the instrument so that the magnesium and carbon lines are recorded on its detectors. The solution is to get a light source that can produce these same lines in the laboratory," he says, and use them to properly adjust the instrument's sensors.



Readily available lamps can simulate the UV light from the singly ionized magnesium, but generating the UV light from triply ionized carbon (carbon with three electrons removed) is difficult. Enter NIST's unique "sliding spark source." It consists of a pair of graphite electrodes with a quartz surface in between. A spark from these electrodes glides along the quartz surface, controllably producing the desired wavelengths of <u>UV light</u> from ionized carbon. Inside a clean room in Huntsville, UV radiation from the spark source entered SUMI, enabling its sensors to be accurately positioned before deployment.

On July 30, 2010, SUMI was successfully launched from White Sands, N.M. It rocketed 320 km in space and observed sunspot 11092 for about 6 minutes before parachuting back to earth. The Huntsville team is analyzing the data it obtained.

Provided by National Institute of Standards and Technology

Citation: NIST ultraviolet source helps NASA spacecraft measure the origins of space weather (2010, September 1) retrieved 3 May 2024 from <u>https://phys.org/news/2010-09-nist-ultraviolet-source-nasa-spacecraft.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.