

Red light regulates nectar secretion

September 27 2010



Ants of the genus *Crematogaster* defend a Lima bean plant (*Phaseolus lunatus*) against an herbivorous insect. Photo: Christian Kost, MPI Chemical Ecology

Flowering plants produce nectar to attract insect pollinators. Some plant species, such as Lima bean, also secrete nectar from so-called extrafloral nectaries to attract ants which in turn fend off herbivores. Scientists of the Max Planck Institute in Jena, Germany, have discovered that the production of extrafloral nectar is light dependent. They have shown that the plants are able not only to distinguish between day and night, but also to adapt their nectar secretion to current light conditions by using a special photoreceptor, the phytochrome.

Phytochrome probably influences the regulation of a special enzyme that binds the [plant hormone](#) jasmonic acid (JA) to the amino acid isoleucine (Ile). The emerging JA-Ile molecule affects the secretion of extrafloral nectar in such a way that the plant's defense against herbivores is most

effective whenever herbivory is most likely - or, more precisely, during the day. (*PNAS* Early Edition)

Plants have to continuously defend themselves against herbivores to survive and reproduce. They do this directly by producing toxic substances, such as nicotine, or indirectly, by calling their enemies' enemies for help. An example of an indirect defense is the release of volatile substances that attract predatory [insects](#) or parasitoids and guide them to their prey; for example, predatory wasps or bugs are led to a caterpillar that is feeding on a plant.

Another indirect defense is the secretion of extrafloral nectar from special leaf organs. In this way Lima beans attract ants that not only enjoy the sweet nectar but also defend the plant against herbivores. Scientists in the Department of Bioorganic Chemistry study this "sweet" defense mechanism. Radhika Venkatesan, a PhD student from India, completed a series of experiments on this topic and tested whether nectar secretion in Lima beans is light dependent. "After all, nectar consists mainly of sugars, and sugars are primary products in the process of photosynthesis - which depends on light," notes the scientist. In the course of her studies, which were published in the Early Edition of the "Proceedings of the National Academy of Sciences USA" last week, she came across an old acquaintance in plant research - the so-called phytochrome. Plants contain phytochromes as photoreceptors in their leaves, which is the reason why phytochromes are sometimes called "the eye of the plant."

"Not the light intensity, but light quality or composition plays a decisive role in regulating nectar production in Lima bean," says Wilhelm Boland, director of the Max Planck Institute. As he explains, the phytochrome in the plant absorbs [red light](#) that enables the plant to distinguish the diurnal and seasonal variation of sunlight quality. Radhika Venkatesan's experiments are the first to demonstrate that

plants also use the phytochrome system to set up their lines of defense effectively and economically.

The hormone jasmonic acid is known as an important signal that plants produce after wounding by herbivores. It also plays a central role in regulating nectar secretion. The scientists have discovered that phytochrome-mediated light regulation has a significant impact on the signaling effect of jasmonic acid: Free jasmonic acid inhibits nectar secretion in the dark but stimulates its production in the light. Radhika Venkatesan found the key to this light-regulated behavior in a reaction that binds jasmonic acid to the amino acid isoleucine. The emerging conjugate JA-Ile is a signal molecule already known from other studies. For the first time, it has been identified as the actual elicitor of nectar secretion. Additional experiments have confirmed that nectar production doesn't increase if binding of JA and isoleucine is prohibited by an inhibitor. If plants are wounded in the dark to stimulate JA production, JA-Ile is produced only in those leaves that were previously exposed to red light.

More information: Venkatesan Radhika, Christian Kost, Axel Mithöfer, Wilhelm Boland (2010). Regulation of extrafloral nectar secretion by jasmonates in lima bean is light dependent. *Proceedings of the National Academy of Sciences USA*, Early Edition, 20. September 2010, [DOI: 10.1073/pnas.1009007107](https://doi.org/10.1073/pnas.1009007107)

Provided by Max Planck Institute

Citation: Red light regulates nectar secretion (2010, September 27) retrieved 9 April 2024 from <https://phys.org/news/2010-09-red-nectar-secretion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.