

GM safety debate may have new twist

October 28 2010



In planta transformation. Image: By Kirsty Foster

By studying plant-fungi-bacteria interactions at plant wound sites, the team have identified a natural process stimulated by a hormone released by the wounded plant that would allow synthetic genes to move across organisms and out into the wild.

The bacterium *Agrobacterium tumefaciens* transforms plant tissue as part of its infection process. This natural process provides an important toolbox for scientists to genetically manipulate many species of plants. Recently this technology has been developed for non-plant organisms including <u>fungi</u> by the Bailey & Foster Group in Bristol's School of Biological Sciences.

Their success has come from adding the plant wound hormone acetosyringone, which triggers *Agrobacterium* transformation mechanisms and allows foreign genes to modify cells (genetic transformation). In the natural environment *Agrobacterium* and fungi



likely encounter each other at plant wound sites where acetosyringone is present, raising the possibility of natural gene transfer from bacterium to fungus.

Professor Gary Foster and colleagues tested whether transformation of fungi by *Agrobacterium* can occur in nature on plants. Their results clearly demonstrate that when placed together on damaged plant tissue, *Agrobacterium* readily transforms associated fungi. "This suggests a previously unknown route for horizontal gene transfer in nature," said Professor Foster.

These results may have implications for the risk assessment of GM plants generated via *Agrobacterium*-mediated transformation. *Agrobacterium* can survive within <u>plant tissue</u> following artificial transformation in tissue culture, and can be detected within regenerated transgenic plants. This research shows that these <u>bacteria</u> have the potential to move the same genetic modifications to fungi in a natural environment.

Prior to release of a GM plant, elimination of *Agrobacterium* following modern genetic modification is a key concern of geneticists and policy makers as it is essential to prevent later escape of synthesised gene from *Agrobacterium* to other organisms.

Professor Foster said: "This study suggests that the encounter between *Agrobacterium* and a fungus on the plant surface may lead to gene flow in a previously overlooked way, potentially leaking GM genes into the natural world."

More information: The Bristol study, published online in *PLoS ONE*, was carried out with financial support from NERC.



Provided by University of Bristol

Citation: GM safety debate may have new twist (2010, October 28) retrieved 3 May 2024 from <u>https://phys.org/news/2010-10-gm-safety-debate.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.